

THE AMERICAN MATHEMATICAL MONTHLY	MAA
The Lattice-Point Area of a Quadratic Form	771
Elliptic Curves, Integer Compositions, and Nets of Antiprisms	786
An Explicit Last Equation	802
Combinatorial Properties of Matrix Chain Inequalities	809
On a Problem of Erdős and Moser	816
On a Problem of Erdős and Moser	825
On a Problem of Erdős and Moser	835
On a Problem of Erdős and Moser	841
On a Problem of Erdős and Moser	848
On a Problem of Erdős and Moser	850
On a Problem of Erdős and Moser	855

The American Mathematical Monthly

ISSN: 0002-9890 (Print) 1930-0972 (Online) Journal homepage: <https://www.tandfonline.com/loi/uamm20>

A Note on the Stammer Hyperbola

Franz Embacher & Hans Humenberger

To cite this article: Franz Embacher & Hans Humenberger (2019) A Note on the Stammer Hyperbola, The American Mathematical Monthly, 126:9, 841-844, DOI: [10.1080/00029890.2019.1644125](https://doi.org/10.1080/00029890.2019.1644125)

To link to this article: <https://doi.org/10.1080/00029890.2019.1644125>



© 2019 The Authors. Published with license by Taylor & Francis Group, LLC.



Published online: 23 Oct 2019.



Submit your article to this journal [↗](#)



Article views: 274



View related articles [↗](#)



View Crossmark data [↗](#)

A Note on the Stammler Hyperbola

Franz Embacher and Hans Humenberger

OPEN ACCESS

Abstract. An alternative characterization of the Stammler hyperbola of a triangle is given.

The Stammler hyperbola of a scalene triangle is the unique conic section on which the incenter, the three excenters, and the circumcenter lie [2–4]. (In case of an isosceles but nonequilateral triangle the hyperbola collapses to a pair of straight lines.) In this note, we show how the Stammler hyperbola appears in a different context.

An arbitrary interior point P of an acute or rectangular triangle $\triangle ABC$ gives rise to a partition of the triangle into six pieces, separated by the line segments from P to the vertices and the perpendiculars from P to the sides. We will call this a P -induced partition of the triangle.




Figure 1. P is an arbitrary interior point of the acute triangle $\triangle ABC$, inducing a partition of the triangle into six pieces. The two groups of alternate pieces are distinguished by different shading. The true trilinear coordinates of P —i.e., its normal distances to the triangle sides—are denoted by ξ , η , and ζ . The triangle sides are divided into two subsegments each, their lengths being denoted by r , r' , s , s' , t , and t' . The angles of the triangle are denoted by α , β , and γ .

As illustrated in Figure 1, the six pieces of the original triangle fall into two groups (distinguished by different shading), each group consisting of three subtriangles sharing no common line segment. We may then ask for which locations of P the sums of the areas of the pieces of each group are equal. We will call the P -induced partition *fair* if this is the case.

doi.org/10.1080/00029890.2019.1644125

MSC: Primary 51M04

Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/uamm.

© 2019 The Authors. Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits noncommercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

Theorem 1. *The P -induced partition of an acute or right scalene triangle is fair if and only if P lies on the Stammler hyperbola of that triangle.*

Proof. We use the notation shown in Figure 1. The condition for the partition to be fair obviously reads

$$\frac{1}{2} r \xi + \frac{1}{2} s \eta + \frac{1}{2} t \zeta = \frac{1}{2} r' \xi + \frac{1}{2} s' \eta + \frac{1}{2} t' \zeta, \quad (1)$$

which may be rewritten as

$$(r - r') \xi + (s - s') \eta + (t - t') \zeta = 0. \quad (2)$$

Some elementary trigonometry reveals that

$$r = \frac{\zeta + \xi \cos \beta}{\sin \beta}, \quad (3)$$

$$r' = \frac{\eta + \xi \cos \gamma}{\sin \gamma}, \quad (4)$$

and three similar expressions for s , s' , t , and t' obtained by cyclic permutation of the respective quantities. Inserting these into (2), we first collect all expressions containing only the angle α :

$$-s' \eta + t \zeta = -\frac{\zeta \eta + \eta^2 \cos \alpha}{\sin \alpha} + \frac{\eta \zeta + \zeta^2 \cos \alpha}{\sin \alpha} = (\zeta^2 - \eta^2) \cot \alpha. \quad (5)$$

The contributions $r \xi - t' \zeta$ and $-r' \xi + s \eta$ now follow simply by cyclic permutation. Collecting everything, the fairness condition (2) takes the form

$$(\cot \beta - \cot \gamma) \xi^2 + (\cot \gamma - \cot \alpha) \eta^2 + (\cot \alpha - \cot \beta) \zeta^2 = 0. \quad (6)$$

This is the equation of a conic section in trilinear coordinates (see [5, pp 154–158] or [6]). In order to identify it, we allow exterior locations of P as well, i.e., trilinear coordinates $\xi : \eta : \zeta$ that are not necessarily all positive. Four points lying on this conic section are readily found:

- (i) the incenter I of the triangle whose trilinear coordinates are $1 : 1 : 1$;
- (ii) the excenters of the triangle whose trilinear coordinates are $-1 : 1 : 1$, $1 : -1 : 1$, and $1 : 1 : -1$, respectively.

All these points satisfy $\xi^2 = \eta^2 = \zeta^2$, thus rendering the left-hand-side of (6) identically zero. A fifth point meeting the condition (6) is the circumcenter O of the triangle. Its trilinear coordinates are given by [1, 6]

$$\cos \alpha : \cos \beta : \cos \gamma. \quad (7)$$

Inserting them into (6) and using $\alpha + \beta + \gamma = \pi$ shows that (6) also holds in this case. Since five distinct points uniquely determine a conic section, (6) is the equation of the Stammler hyperbola.

Thus we have shown that the set of all points P inducing a fair partition of the triangle are exactly those points on the Stammler hyperbola that lie in the interior of the given triangle. ■




Figure 2. The P -induced partition of the acute scalene triangle $\triangle ABC$ is fair if and only if P lies on the Stammler hyperbola.

In Figure 2 a configuration with P on the Stammler hyperbola is shown.

Remark (Generalization to arbitrary scalene triangles). So far, we have assumed the triangle $\triangle ABC$ to be acute or rectangular. Allowing $\triangle ABC$ to have an obtuse angle, we face the problem that one of the perpendiculars from P to the side lines may partially fall outside the triangle. This spoils the beautiful concept that every interior point P gives rise to a partition of the triangle into six pieces as shown in Figure 1. A simple possibility to generalize the previous result to arbitrary scalene triangles is simply to exclude all interior points for which that problem occurs. Hence, we call an interior point P *admissible* if all three perpendiculars from P to the side lines lie inside the triangle. Then

- (i) any admissible point P induces a partition of the triangle, and
- (ii) if this partition is fair, then P lies on the Stammler hyperbola of that triangle.

In order to include the reverse direction we must take into account that if the triangle has an obtuse angle the Stammler hyperbola contains nonadmissible interior points. Hence

- (iii) an admissible point P induces a fair partition if and only if it lies on the (obviously defined) admissible part of the Stammler hyperbola.

Another way of generalizing our alternative characterization of the Stammler hyperbola to arbitrary scalene triangles—on a formal level—is to retain (3), (4), and the analogous expressions for s , s' , t , and t' for every interior point P . However, since one of these quantities may become negative, they should be interpreted as *oriented* lengths. As a consequence, the fairness condition (1) may be interpreted as a balance equation in terms of *oriented* areas instead of areas. If, for example, the angle γ is obtuse, and P is chosen such that $r' < 0$ (see Figure 3), then the quantity $\frac{1}{2}r'\xi$ in (1) is negative. It represents the oriented area of the rectangular triangle $\triangle PCD$, where D is the intersection of the perpendicular from P to BC with BC (i.e., the pedal point corresponding to the trilinear coordinate ξ). All the other five terms in (1) correspond to subtriangles of $\triangle ABC$ and contribute positive areas. Since D lies outside of $\triangle ABC$, the triangle $\triangle PCD$ is not a subtriangle of $\triangle ABC$. Thus, although in such a case the




Figure 3. If the angle γ is obtuse, and P is chosen such that $r' < 0$ then the dotted area, represented by the quantity $\frac{1}{2}r'\xi$, counts as negative. In such a case the fairness condition (1) still holds in a formal sense, but it is no longer associated with a partition of $\triangle ABC$.

fairness condition is rescued in a formal sense and still leads to the Stammler hyperbola, it is no longer associated with a partition of $\triangle ABC$. In much the same way, this scheme may be generalized to points P outside the original triangle (in which case one or two of the trilinear coordinates ξ , η , and ζ are nonnegative), leading to a characterization of its complete Stammler hyperbola.

REFERENCES

- [1] Kimberling, C. (1994). Central points and central lines in the plane of a triangle. *Math. Mag.* 67(3): 163–187.
- [2] Kimberling, C. (2018). Encyclopedia of triangle centers. faculty.evansville.edu/ck6/encyclopedia/ETC.html
- [3] Phelps, S. (2018). The Stammler hyperbola. geogebra.org/m/yUNsGaZV#material/rSgkR6MP
- [4] Weisstein, E. W. (2018). Stammler Hyperbola. From *MathWorld*—A Wolfram Web Resource. mathworld.wolfram.com/StammlerHyperbola.html
- [5] Whitworth W. A. (1866). *Trilinear Coordinates and Other Methods of Modern Analytical Geometry of Two Dimensions: An Elementary Treatise*. Cambridge, UK: Deighton, Bell, and Co.
- [6] Wikipedia. (2018). Trilinear coordinates. en.wikipedia.org/wiki/Trilinear_coordinates

Faculty of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, A-1090 Wien, Austria
franz.embacher@univie.ac.at

Faculty of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, A-1090 Wien, Austria
hans.humenberger@univie.ac.at