

Cytotype stability, facultative apomixis and geographical parthenogenesis in *Ranunculus kuepferi* (Ranunculaceae)

Anne-Caroline Cosendai and Elvira Hörandl*

Department of Systematic and Evolutionary Botany, Faculty Centre for Biodiversity, University of Vienna, Rennweg 14, 1030 Vienna, Austria

*For correspondence. E-mail elvira.hoerandl@univie.ac.at

Received: 24 August 2009 Returned for revision: 12 October 2009 Accepted: 23 November 2009 Published electronically: 24 January 2010

- **Background and Aims** Asexual organisms are more widespread in previously glaciated areas than their sexual relatives ('geographical parthenogenesis'). In plants, this pattern is probably dependent on reproductive isolation and stability of cytotypes within their respective distribution areas. Both partial apomixis and introgressive hybridization potentially destabilize the spatial separation of sexual and apomictic populations. The wide distribution of apomicts may be further enhanced by uniparental reproduction which is advantageous for colonization. These factors are studied in the alpine species *Ranunculus kuepferi*.
- **Methods** Geographical distribution, diversity and mode of reproduction of cytotypes were assessed using flow cytometry and flow cytometric seed screening on samples from 59 natural populations of *Ranunculus kuepferi*. Seed set of cytotypes was compared in the wild.
- **Key Results** Diploid sexuals are confined to the south-western parts of the Alps, while tetraploid apomicts dominate in previously glaciated and in geographically isolated areas despite a significantly lower fertility. Other cytotypes (3x, 5x and 6x) occur mainly in the sympatric zone, but without establishing populations. The tetraploids are predominantly apomictic, but also show a partial apomixis via an uncoupling of apomeiosis and parthenogenesis in the seed material. Both pseudogamy and autonomous endosperm formation are observed which may enhance uniparental reproduction.
- **Conclusions** Diploids occupy a glacial relic area and resist introgression of apomixis, probably because of a significantly higher seed set. Among the polyploids, only apomictic tetraploids form stable populations; the other cytotypes arising from partial apomixis fail to establish, probably because of minority cytotype disadvantages. Tetraploid apomicts colonize previously devastated and also distant areas via long-distance dispersal, confirming Baker's law of an advantage of uniparental reproduction. It is concluded that stability of cytotypes and of modes of reproduction are important factors for establishing a pattern of geographical parthenogenesis.

Key words: Apomixis, flow cytometry, geographical parthenogenesis, glaciations, polyploidy, *Ranunculus kuepferi*.

INTRODUCTION

Vandel (1928) presented the term 'geographical parthenogenesis' for the phenomenon that related sexual and asexual taxa have different distribution areas. Later, several authors discussed the causality of the pattern for animals and plants by addressing a number of new aspects related to the wider distributions of asexuals (Bell, 1982; Bierzychudek, 1985; Asker and Jerling, 1992; Law and Crespi, 2002; Van Dijk, 2003; Kearney, 2005, 2006; Hörandl, 2006; Lundmark, 2006). Plants reproducing via apomixis, i.e. via asexually formed seed, tend to grow at higher altitudes and latitudes, and colonize more frequently previously glaciated areas than their sexual relatives (Bierzychudek, 1985; Van Dijk, 2003; Hörandl *et al.*, 2008).

Asexual reproduction is in general frequently connected to polyploidy, and almost all apomictic plants are polyploid (Asker and Jerling, 1992). The tight connection of polyploidy and apomixis usually leads to differentiation and reproductive isolation of cytotypes (Van Dijk, 2007; Kao, 2007; Mráz *et al.*,

2008). However, facultative apomixis can increase considerably the cytotype diversity within apomictic populations. Apomixis usually requires the co-ordination of embryo sac development without meiosis (apomeiosis) and the development of the egg cell without fertilization (parthenogenesis). The uncoupling of these processes leads to shifts in ploidy levels (Nogler, 1984a); meiosis plus parthenogenesis results in a dihaploid offspring ($n + 0$), while fertilization of an apomeiotic egg cell results in an increase of ploidy level ($2n + n$, BIII hybrids). Dihaploids and BIII hybrids are expected to undergo a decrease in fitness (Van Dijk and Vijverberg, 2005); continued reduction of ploidy levels results in the expression of previously masked recessive disadvantageous alleles, while continued increase in ploidy levels is limited by cellular constraints and functional disturbances of regulation mechanisms (Comai, 2005). A newly arisen, partially sexual cytotype may further suffer from minority cytotype disadvantages in the population (Levin, 1975), because it will mostly receive pollen of the wrong ploidy level. This may

not only have negative effects on the fitness of the embryo, but also on endosperm formation. In the endosperm of flowering plants, a ratio of two maternal and one paternal copies of the genome are optimal for development, probably because of genomic imprinting; deviations from this ratio are sometimes tolerated, but often result in disturbances in seed formation (Vinkenoog *et al.*, 2003, Spielmann *et al.*, 2003; Talent, 2009). Since the majority of apomictic plants are pseudogamous and need pollen for endosperm fertilization, interploidal crosses pose a problem because they cause endosperm imbalance and potentially seed abortion. For these reasons, stability of cytotypes is probably an important factor for fitness and the distributional success of apomictic lineages. In fact, patterns of geographical parthenogenesis have so far not been reported for those taxa with a highly facultative and unstable apomixis (reviewed in Hörandl *et al.*, 2008).

Most apomicts produce fertile, meiotically reduced pollen, and the genetic factors controlling apomixis can be inherited via the pollen (e.g. Asker and Jerling, 1992; Mogie, 1992). In mixed populations, an apomictic pollen donor can fertilize a sexual plant, thereby transferring apomixis to the offspring of the sexual. Under a simple model of dominant, single locus control for apomixis and an apomictic pollen donor heterozygous for the apomixis factor, some of the offspring of the sexual will become apomictic. In turn, the pollen of the sexual does not fertilize an apomictic plant, because the egg cell develops parthenogenetically. This unidirectional hybridization would result in introgression of apomixis from the apomicts into the sexual populations, but not vice versa. Theoretically, sexuality should disappear from the population after a few generations (Mogie, 1992; Mogie *et al.*, 2007). Among other factors, this process could contribute to geographical parthenogenesis by replacing sexuals by apomicts in sympatric areas. However, Mogie (1992) has already pointed out that the amount of actual introgression also depends on female fertility, and a significantly higher fertility of sexuals prevents their replacement by apomictic cytotypes (Hörandl and Tensch, 2009). The fertility of cytotypes is therefore an important factor for geographical parthenogenesis.

Other theories explain geographical parthenogenesis by superior colonizing abilities. The capacity to found a new population from a single individual or seed is a big advantage for colonization, especially after long-distance dispersal (Baker's law; Baker, 1967; Hörandl, 2008, Hörandl *et al.*, 2008). Here apomixis with autonomous endosperm formation provides an advantage over pseudogamy which still needs pollination, unless the plants can use self-pollen for endosperm fertilization (Dickinson *et al.*, 2007; Hörandl, 2008). Sexuals, in contrast, are probably more efficient in habitats with regular pollinator frequencies and benefit from the advantage of genetic diversity. Since sexual species of apomictic complexes are usually self-sterile (Dickinson *et al.*, 2007; Hörandl, 2009b), their ability to found populations in geographically distant areas is limited by the need of mating partners and pollinators.

Other theories explaining patterns of geographical parthenogenesis rely rather on genetic diversity of populations and a different response of sexual and apomictic populations to variable environments [e.g. host–parasite interactions (Van Valen, 1973; Vorburger, 2006); the benefit of general purpose genotypes (Lynch, 1984); niche differentiation of clones (Vrijenhoek, 1984, 1994)]. However, genetic diversity and

the response to selection can be altered by polyploidy (Levin, 2002) and by clonal diversity (e.g. Van Dijk, 2003). Therefore, the assessment of distribution and stability of cytotypes and of modes of reproduction is indirectly of crucial importance for these models.

The alpine species *Ranunculus kuepferi* is an interesting model system for studying patterns and processes of geographical parthenogenesis in previously glaciated areas. Küpfer (1974) first recognized it as a separate species with diploid ($2n = 16$) and tetraploid ($2n = 32$) cytotypes. He found diploids only in the south-western parts of the Alps that have remained ice-free during the last glacial maximum of Würm glaciation (approx. 10 000 years ago). This area has long been known to be a glacial refugium for many plant species (Merxmüller, 1952, 1953, 1954; Schönswetter *et al.*, 2005). The tetraploids, in contrast, were observed in the previously glaciated central western Alps. Later, Huber (1988) and Burnier *et al.* (2009) refined the distribution and reported the tetraploid cytotype eastwards to Eastern Tyrol (Austria). They assessed the presence of triploids ($2n = 24$) in the sympatric area of the diploid and the tetraploid cytotype, suggesting a hybrid zone. Outside the Alps, tetraploid cytotypes have been detected in Corsica and in the Apennines (Küpfer, 1974; Huber, 1989; Burnier *et al.*, 2009).

Küpfer (1974) had already observed reduced pollen fertility (50–80% aborted pollen grains), and reduced fertility (10–100% of achenes aborted) in the tetraploid cytotype, while the diploid cytotypes had good pollen (0–20% aborted) and achenes (0–10% aborted). Information on other cytotypes and statistical evaluations of differences, however, were missing so far. An embryological observation by Burnier *et al.* (2009) on a single tetraploid specimen suggested an embryo-sac formation similar to the well-studied apomictic model system *R. auricomus* (Nogler, 1984a, b, 1995). Here meiosis takes place, but the megaspore tetrad aborts during the later development. Instead, a somatic cell of the nucellus develops into an unreduced, 8-nucleate embryo sac of the *Polygonum* type. This apomeiotic (aposporous) embryo sac development is coupled to a parthenogenetic development of the egg cell. These two processes are facultative and can be uncoupled, resulting in shifts of ploidy levels in the offspring (Nogler, 1984a, b, 1995).

In *R. kuepferi*, mode of reproduction and cytotype diversity within populations has not yet been assessed on population samples throughout the distributional range. Furthermore, the ploidy level and mode of reproduction of some geographically isolated outposts in the North Apennines (Mt Cusna) and in the eastern central Alps (populations around Turracherhöhe) were so far unknown. These areas potentially could represent glacial refugia for sexual diploid populations. The easternmost populations (Fig. 1, no. 59) are located very near to the last glacial maximum eastern refugia, a spot well known for its endemic flora (Tribsch, 2004; Schönswetter *et al.*, 2005). Disjunct peripheral distributions of diploid species with related polyploid cytotypes in the centre of the Alps have been observed, for example, in the *R. auricomus* complex (Paun *et al.*, 2006; Hörandl, 2009a) and in sexual *Biscutella laevigata* (Parisod and Besnard, 2007). In North America, diploid sexual cytotypes of *Townsendia hookeri* have a strongly disjunct distribution in peripheral refugia of the Wisconsin glaciation, while polyploid apomicts occur in the central previously glaciated area (Thompson and Whitton, 2006). Alternatively,




FIG. 1. Map of the distribution of *Ranunculus kuepferi*. Small black squares connected to the pie diagrams indicate the location, population numbers correspond to Table 1. Pie diagrams present the proportions of cytotypes for each population, as indicated. The black line indicates the extension of the last glacial maximum of the Würm glaciation.

these outposts in *R. kuepferi* could have been founded after long-distance dispersal by tetraploid apomicts. This scenario would support an idea of superior founder abilities of tetraploid apomicts according to Baker's law.

For *R. kuepferi*, a detailed study on cytotype diversity and mode of reproduction throughout the range of the species has been missing so far, i.e. it is uncertain whether the species shows a pattern of geographical polyploidy or geographical parthenogenesis. In the light of the reduced fertility of tetraploid cytotypes, their distributional success appears to be a paradox; advantages of apomixis may explain the observed pattern. Therefore, it is desirable to study stability and modes of apomixis of the species throughout its distributional range and to test stability of cytotypes and modes of reproduction within the respective areas. For the latter, possible introgression of apomixis into sexual species within the sympatric area, and the female fertility of cytotypes are of interest. It is followed by testing whether geographically isolated outposts outside the ice-shield of the Würm glaciation represent diploid sexual refugia or colonies of tetraploid apomicts. For these investigations into cytotype diversity, involving large sample sizes, the use of flow cytometric methods is a powerful approach. To assess modes of reproduction and pathways of embryo formation, flow cytometric seed screening

(FCSS) is a highly efficient method (Matzk *et al.*, 2000). The focus here is on intrinsic features of apomixis as factors of geographical parthenogenesis; patterns of genetic diversity within and among populations, and hypotheses on origins of apomictic cytotypes will be presented in forthcoming papers.

The following specific questions were addressed in the study. Does this distribution pattern represent a geographic distribution of sexual polyploidy or geographical parthenogenesis? How constant is the ploidy level within populations over all the distribution range, and how can new cytotypes be formed? In the hybrid zone, is there an indication of introgression of apomixis into the sexual populations, and do apomicts have a potential to replace the sexual lineages? Are the areas outside or near the margin of the previous ice-shield refugia of diploid sexual populations or is a colonization scenario by tetraploid apomicts more likely?

MATERIALS AND METHODS

Plant material

The plants of the species used in this research, *Ranunculus kuepferi* Greuter et Burdet, were collected in the wild during spring and summer 2004–2007; for details, see Table 1 and

TABLE 1. Provenance of materials used in this study

Population no.*	Country†	Province	Locality‡	Altitude	Latitude	Longitude	Collector§	Date
1	F	Corse-du-Sud	Corsica	1541 m	42°01'46.4"	9°12'34.5"	ACC	28 May 2005
2	I	Emilia-Romagna	Mt Cusna	1594 m	44°18'06.7"	10°22'26.6"	ACC-AC	26 May 2006
3	F	Var	La Chens I	1610 m	43°44'59.3"	6°39'25.5"	PK	4 June 2004
4	F	Var	La Chens II	1607 m	43°44'58.5"	6°39'29.1"	ACC-AC	28 May 2006
5	F	Alpes Maritimes	Col de Tende	1888 m	44°09'03.0"	7°33'56.3"	ACC	9 June 2004
6	I	Piemonte	Valle di Pesio I 9589	1700 m	44°11'43.68"	7°39'33.85"	EH	9 July 2007
7	I	Piemonte	Valle di Pesio II 9592	1700 m	44°11'43.68"	7°39'33.85"	EH	10 July 2007
8	I	Piemonte	Passo del Duca 9525/9534	1700 m	44°11'43.68"	7°39'33.85"	EH	14 July 2004
9	I	Piemonte	Valle di Pesio III 9593	1925 m	44°11'43.68"	7°39'33.85"	EH	10 July 2007
10	I	Piemonte	Vallone Cravina 9595	1960 m	44°13'24.72"	7°37'11.16"	EH	12 July 2007
11	I	Piemonte	Col della Perla I 9596	2080 m	44°9'11.05"	7°37'21.14"	EH	13 July 2007
12	I	Piemonte	Col della Perla II 9597	2200 m	44°9'11.05"	7°37'21.14"	EH	14 July 2007
13	F	Alpes Maritimes	Notre Dame de la Fenestre	1885 m	44°05'45.3"	7°21'34.2"	ACC	11 June 2004
14	F	Alpes Maritimes	Col d'Isola	2210 m	44°11'43.7"	7°09'19.7"	ACC	11 June 2004
15	I	Piemonte	Colle della Lombarde I 9601	2260 m	44°12'25.68"	7°8'51.98"	EH	15 July 2007
16	I	Piemonte	Colle della Lombarde II 9602	2477 m	44°13'17.60"	7°9'9.25"	EH	16 July 2007
17	F	Alpes de Haute Provence	Vallette	1820 m	44°07.5'	6°33.5'	PK	4 June 2004
18	F	Alpes de Haute Provence	Champs I	1925 m	44°09'59.5"	6°42'28.0"	ACC	13 June 2004
19	F	Alpes Maritimes	Champs II	2080 m	44°10'33.8"	6°41'53.0"	ACC-AC	29 May 2006
20	F	Alpes Maritimes	Cayolle I	2193 m	44°15'13.6"	6°44'52.1"	ACC	13 June 2004
21	F	Alpes de Haute Provence	Cayolle II	2325 m	44°15'34.4"	6°44'41.3"	ACC-AC	30 May 2006
22	F	Alpes de Haute Provence	Allos I	2247 m	44°22"	6°37"	PK	4 June 2004
23	F	Alpes de Haute Provence	Allos II	2080 m	44°18'03.4"	6°35'06.0"	ACC-AC	29 May 2006
24	F	Hautes Alpes	Vars	2134 m	44°32'21.3"	6°42'05.6"	ACC	14 June 2004
25	F	Hautes Alpes	Raboux	1859 m	44°38'38.4"	5°58'43.1"	ACC	15 June 2004
26	F	Hautes Alpes	Haute Queyras, Col de La Croix	2000 m	44°46'0"	7°01"	MH	22 May 2004
27	F	Hautes Alpes	Haute Queyras, Montette	2000 m	44°50'0"	6°55'0"	MH	23 May 2004
28	F	Drôme	Vercors I	1470 m	44°54'07"	5°28'039"	PK	3 June 2004
29	F	Drôme	Vercors II	1325 m	44°50'26.1"	5°25'23.1"	ACC-AC	30 May 2006
30	F	Hautes Alpes	Col de Lautaret	2060 m	45°02'40.6"	6°24'04.1"	ACC	15 June 2004
31	F	Savoie	Mt Cenis	2025 m	45°13'55.4"	6°53'53.6"	ACC-AC	3 July 2007
32	F	Savoie	Col d'Iseran	2768 m	45°25'09.2"	7°01'52.9"	ACC-AC	3 July 2007
33	I	Valle d'Aoste	Gran Paradiso	2079 m	45°37'0.5"	7°33'12.2"	ACC-AC	4 July 2007
34	F	Savoie	Petit St Bernard	2215 m	45°40'40.4"	6°52'55.2"	ACC-AC	2 July 2007
35	CH	Valais	Grand St Bernard	2380 m	45°52'09.9"	7°09'33.3"	ACC-AC	2 July 2007
36	CH	Valais	Arpilles	1830 m	46°04.94'	7°.78'	AC	18 June 2006
37	CH	Valais	Ovronnaz	1840 m	46°13.03'	7°09.52'	AC	18 June 2006
38	I	Valle d'Aoste	Cervinia	2200 m	45°55'54.6"	7°38'18.1"	ACC-AC	4 July 2007
39	CH	Valais	Jeizinen	2020 m	46°20'07"	7°43'85"	ACC	5 July 2004
40	CH	Valais	Lötschental	1773 m	46°26'05.5"	7°51'48.0"	ACC	29 June 2006
41	CH	Valais	Simplon Pass	2019 m	46°15'03.2"	8°01'48.2"	ACC	29 June 2006
42	CH	Valais	Furka Pass	2162 m	46°34'45.8"	8°25'30.9"	ACC	28 June 2006
43	CH	Ticino	Lukmanier Pass	1946 m	46°33'49.2"	8°47'54.7"	ACC	28 June 2006
44	CH	Graubunden	Rheinwald 9603	2100 m	46°33'16.05"	9°14'52.32"	EH	18 July 2007
45	CH	Graubunden	Julier Pass	2277 m	46°28'20.9"	9°44'01.4"	ACC	27 June 2006
46	CH	Graubunden	Albula Pass	2312 m	46.58333°	9.83333°	ACC	26 June 2006
47	CH	Graubunden	Bernina Pass	2301 m	46°24'47.3"	10°01'17.3"	ACC	27 June 2006
48	A	Vorarlberg	Arlberg Pass	2269 m	47°08'49.1"	10°14'55.3"	ACC	25 June 2006
49	CH	Graubunden	Umbrail Pass	2463 m	46°32'51.3"	10°26'06.3"	ACC	26 June 2006
50	A	Tirol	Kaunertal	2525 m	46°52'21.8"	10°42'37.5"	ACC	25 June 2006
51	I	Trento	Tonale Pass	2400 m	46°16'21.27"	10°34'40.88"	EH	14 July 2006
52	A	Tirol	Timmelsjoch	2105 m	46°55'13.5"	11°03'10.6"	ACC	24 June 2006
53	A	Tirol	Tuxer Alps	2315 m	47°07'	11°34'	CS	6 July 2006
54	I	Trento	Rosengarten	2500 m	46°27'19.56"	11°37'56.51"	EH	11 July 2006
55	I	Trento	Padon Pass	2350 m	46°27'47.80"	11°53'42.88"	EH	9 July 2006
56	I	Veneto	Mt Dürrenstein	2400 m	46°39'39.24"	12°10'57.86"	EH	18 July 2006
57	A	Carinthia	Mt Großglockner	2220 m	47°4'47.63"	12°45'50.37"	EH	12 August 2005
58	A	Carinthia	Mt Sadnig	2200 m	46°57'42"	13°35"	PS-GS	7 July 2006

Continued

TABLE 1. *Continued*

Population no.*	Country†	Province	Locality‡	Altitude	Latitude	Longitude	Collector§	Date
59	A	Carinthia	Turracherhöhe	2220 m	46°55'20-47"	13°52'44-35"	EH	14 August 2005

* The numbers correspond to those used in Fig. 1.

† F, France; I, Italy; A, Austria; CH, Switzerland.

‡ A roman number indicates repeated sampling on the same population in different years; arabic numbers are herbaria numbers of E. Hörandl (vouchers were deposited in the herbarium of the University of Vienna, WU).

§ ACC, Anne-Caroline Cosendai; ACC-AC, Anne-Caroline Cosendai and André Cosendai; CS, Christoph Seger; EH, Elvira Hörandl; MH, Marc Hämmerli; PK, Philippe Küpfer; PS-GS, Peter Schönswetter and Gerald Schneeweiss.

Fig. 1. About one-third of the individuals were cultivated in the experimental fields of the botanical garden of the University of Vienna. The rest of the samples were dried in silica gel for further molecular analysis. Mature achenes were collected in the wild and dried with silica gel, or taken from the plants in the experimental garden directly after collection (which means that embryo-sac formation and fertilization already has happened before on the natural site) and stored in the fridge at 4 °C.

Ploidy-level determination

Previous chromosome counts of root tip squashes (Cosendai, 2005) were used to fix genome size measures to ploidy levels. Samples were measured via flow cytometry (FC) on fresh and silica-dried material of leaves. The difference in genome size between dried and fresh material was never >10 % of variation, comparable to results obtained by Suda and Trávníček (2006) for silica-dried material. It is noted that the genome size of the fresh material is, in this case, almost always slightly smaller than the silica-gel material; some of the smallest DNA content values are due to old silica gel-dried material (>4 years old) and look very degraded (yellow brownish material). These samples are indicated with an asterisk in Table 3 and may be unreliable for absolute genome-size estimates, but nevertheless, allow for a reliable assessment of ploidy levels. Nuclei extraction was prepared following the procedure presented in Galbraith *et al.* (1983) and Doležel *et al.* (2007a). Leaf material was chopped in Otto buffer I modified (with 1.26 g citric acid and 6 mL Triton X-100 for 60 mL final volume) (Doležel *et al.*, 2007b; Temsch *et al.*, 2008) with *Pisum sativum* (Ps) line 'Ctirad' but mostly with *Zea mays* (Zm) line 'CE-777' seedlings (4.38 pg and 2.73 pg DNA/2C, respectively; Doležel *et al.*, 1998; Greilhuber, 2008) as standard; then RNase (3 mg mL⁻¹; Sigma) was added to the extract and incubated at 37 °C for 0.5 h in a water bath. Otto II buffer with propidium iodide (VWR international; final concentration 50 µg mL) (on the basis of Baranyi and Greilhuber, 1996; Temsch *et al.*, 2009) was added to the suspension of nuclei and stored at 4 °C for about 1 h. Samples were analysed on a CyFlow ML flow cytometer (Partec, Muenster, Germany) equipped with a green laser (100 mW, 532 nm, Cobolt Samba; Cobolt AB, Stockholm, Sweden). Five thousand particles were measured per run and mostly three runs were conducted per sample. Analyses of the runs and peak detection

were made with the FloMax[®] software 2.0.0.1 (Partec, Muenster, Germany). The Cx value in the sense of Greilhuber *et al.* (2005) was calculated based on a linear relationship between the standard and the sample fluorescence intensity. The mean values are calculated on the basis of measuring 25 samples per populations. Preparations were mostly repeated three times for statistical stability. Several measures were done by pooling the samples of several individuals. Appearance of a single peak indicated that the pooled samples were of the same genome size and consequently of the same ploidy level. The picogram and Cx values were calculated per population (Table 2).

Low differences in standard deviations and low coefficients of variation (mostly below 3 %; Table 2) indicate a rather stable genome size within populations, falling clearly into distinct classes corresponding to cytotypes. A regression analysis confirmed that the increase in genome size with ploidy levels is almost perfectly linear, so that ploidy levels are almost exactly double or 1.5-fold the previous ploidy level (Fig. 2). This analysis confirms the reliability of the measurements for the ploidy level assessments.

FCSS

This part of the research is based on mature seeds and the mode of reproduction on the basis of *Polygonum* embryo sac type, as found by Vuille and Küpfer (1985) in the species *Ranunculus parnassifolius* and by Nogler (1984b, 1995) in *Ranunculus auricomus*. Furthermore, some recent embryological observations (Burnier *et al.*, 2009; J. Wagner, pers. comm.) confirm that *Ranunculus kuepferi* has an embryo sac of the *Polygonum* type. The principle here is to measure the quantity of the genome in the embryo and the endosperm, which appear in two different peaks in the histogram file. With this information, one can determine the ploidy level of the embryo and endosperm, and reconstruct the pathway of seed formation (Matzk *et al.*, 2000). Achenes were prepared from 58 individuals according to the same protocol as for leaves. The achenes were softened in Otto I buffer for about 5 min on ice and then chopped following the same procedure as in FC with *Zea mays* (Zm) or *Pisum sativum* (Ps) standard. Per sample, four to seven achenes were used, depending on the quality and the quantity of fruits produced by the plant. Figure 3 illustrates an example of the interpretation of an FCSS histogram. Beside the peaks of the standard (Zm), three peaks for the seeds were observed: the first peak (Rk) is the embryo corresponding to a triploid Cx

TABLE 2. Summary of the population's genome size and ploidy level

Populations names*	Mean values of population (pg)	Standard deviation	Coefficient of variation (%)	No. of individuals	Percentage of ploidy level in the population
2x cytotypes					
Vallone Cravina 9595 (s)	4.433	0.040	0.911	17	100.0
Lachens II (s)	4.144	0.069	1.675	22	100.0
Colle della Lombarde I 9601 (s)	4.459	0.091	2.037	24	100.0
Colle della Lombarde II 9602 (s)	4.405	0.027	0.602	11	100.0
Valle di Pesio I 9589 (s)	4.429	0.036	0.805	25	100.0
Valle di Pesio II 9592 (s)	4.462	0.042	0.946	4	100.0
Valle di Pesio III 9593 (s)	4.419	0.045	1.015	14	100.0
Passo del Duca 9525 (s) [†]	3.820	0.024	0.639	5	100.0
Valette (s) [†]	3.941	0.164	4.149	24	79.2
Vercors II (s)	4.058	0.050	1.220	18	100.0
Mean	4.257	0.244			
3x cytotypes					
Allos I (f)	6.322	0.057	0.909	3	66.7
Allos I (s)	6.487	0.058	0.893	12	91.7
Allos II (s)	6.487	0.058	0.891	24	45.8
Champs I (s)	6.431	0.015	0.228	1	100.0
Champs II (f)	6.269	0.246	3.920	23	47.8
Champs II (s)	6.450	0.040	0.624	7	28.6
Gran Paradiso (s)	5.519			24	4.2
Col della Perla I 9596 (s)	6.599	0.079	1.195	23	47.8
Col della Perla II 9597 (s)	6.652	0.072	1.078	22	27.3
Valette (s) [†]	5.964	0.318	5.325	24	20.8
Mean	6.318	0.340			
4x populations					
Arpilles (s)	8.372	0.223	2.666	19	100.0
Albula (s)	8.704	0.095	1.087	25	100.0
Allos I (s)	8.559	0.021	0.246	12	8.3
Allos II (s)	8.597	0.128	1.485	24	25.0
Allos I (f)	8.015	0.000	0.000	3	33.3
Allos II (f)	8.341	0.025	0.299	2	50.0
Bernina Pass (s)	8.707	0.226	2.599	24	95.8
Cayolle II (f)	8.468	0.135	1.596	20	95.0
Cayolle II (s)	8.543	0.142	1.662	9	88.9
Cervinia (s)	8.578	0.038	0.438	25	100.0
Champs II (f)	8.419	0.082	0.973	23	52.2
Champs II (s)	8.631	0.032	0.376	7	71.4
Corsica (f)	8.196	0.277	3.385	2	100.0
Mt Cusna (f)	8.552	0.017	0.201	4	100.0
Dürrenstein (s)	8.514	0.093	1.097	10	100.0
Furka Pass (s)	8.524	0.071	0.831	25	100.0
Gd Paradiso (s)	8.671	0.361	4.163	24	95.8
Grand St Bernard (s)	8.716	0.091	1.048	24	100.0
Col d'Iseran (s)	8.603	0.049	0.572	25	100.0
Julier Pass (s)	8.251	0.051	0.621	25	100.0
Kaunertal (s)	8.603	0.097	1.132	25	100.0
Lötschental (s)	8.136	0.103	0.025	25	100.0
Lukmanier (s)	8.158	0.059	0.720	25	100.0
Mt Cenis (s)	8.596	0.094	1.098	24	100.0
Ovronnaz (s)	8.552	0.008	0.096	8	100.0
Padon (s)	8.835	0.092	1.044	25	100.0
Col della Perla I 9596 (s)	8.658	0.125	1.438	23	34.8
Col della Perla II 9597 (s)	8.643	0.089	1.027	22	68.2
Pt St Bernard (s)	8.702	0.068	0.776	24	100.0
Rheinwald 9603 (s)	8.547	0.067	0.781	21	100.0
Rosengarten (s)	8.584	0.067	0.780	20	100.0
Mt Sadnig (s)	8.749	0.067	0.771	15	100.0
St Anton (s)	8.834	0.094	1.068	25	100.0
Simplon (s)	8.808	0.247	2.803	25	100.0
Timmelsjoch (f)	8.314	0.077	0.931	1	100.0
Timmelsjoch (s)	8.559	0.146	1.702	25	100.0
Tonale Pass (f)	8.462	0.014	0.160	1	100.0
Turracherhöhe (s)	8.617	0.065	0.753	26	100.0
Tuxer Alps (s)	8.608	0.062	0.724	25	100.0
Umbrail Pass (s)	8.731	0.295	3.375	25	100.0

Continued

TABLE 2. *Continued*

Populations names*	Mean values of population (pg)	Standard deviation	Coefficient of variation (%)	No. of individuals	Percentage of ploidy level in the population
Mean	8.552	0.184			
5x populations					
Allos II (f)	10.337	0.134	1.292	2	50.0
Allos II (s)	10.649	0.163	1.533	24	29.2
Col della Perla I 9596 (s)	10.633	0.121	1.134	23	13.10
Col della Perla II 9597 (s)	10.755	0.052	0.484	22	4.50
Mean	10.540	0.175			
6x populations					
Bernina (s)	12.685	0.220	1.731	24	4.20
Cayolle II (f)	12.671	0.114	0.900	20	5.0
Cayolle II (s)	13.024	0.118	0.905	9	11.10
Perla I 9596 (s)	12.916	0.062	0.482	23	4.30
Mean	12.793	0.199			
Summary of ploidy level	No. of individuals	Percentage of cytotype			
2x	159	15.7			
3x	61	6.02			
4x	778	76.8			
5x	12	1.18			
6x	3	0.3			
Total	1013				

* (s) or (f) at the end of the population name indicates silica-dried or fresh material, respectively.

† Indicates old material (collected in 2004).




FIG. 2. Linear regression of the relative ploidy level Cx value and the genome size in picograms based on leaf material. The slope of the regression is indicated.

value; the second peak is interpreted as the G_2 of the embryo as it has double the amount of DNA of the embryo peak. The third peak has a ratio/picogram value of 10 Cx, corresponding to an endosperm that has arisen from two polar nuclei ($3x + 3x$) plus fertilization of two pollen nuclei ($2x + 2x$). Based on these data, it was possible to determine the formation of seeds and the assumed mode of reproduction.

Fertility of cytotypes

Female fecundity of cytotypes was assessed on 116 individuals from 21 populations, by using achenes collected in the wild or directly after transfer to the experimental garden. The number of well-developed and aborted achenes was




FIG. 3. (A) Histogram of FCSS with five peaks: 1 and 2, standard *Zea mays* in the G_1 and G_2 phase, respectively; 3–5, *Ranunculus kuepferi* (Rk): 3, 3 Cx, embryo G_1 phase; 4, embryo G_2 phase; 5, 10 Cx endosperm (peak G_1 phase). (B) Scheme of the respective embryo sac after fertilization, illustrating a triploid unfertilized embryo corresponding to peak 3, and the 10 Cx endosperm corresponding to peak 5. Since the mother plant was tetraploid, the embryo sac must have developed via a disturbed meiosis. The 3x em developed without fertilization, the endosperm via pseudogamy.

assessed as in Hörandl (2008). Percentages of well-developed achenes per collective fruit were calculated, pooled for cytotypes. After arcsin transformation of percentages, the significance of differences between cytotypes was tested via

one-way ANOVA; for this test, the single sample from a hexaploid plant was pooled with the values of pentaploids. Since Levene's statistic revealed unequal variances among groups, Tanhame's test for pairwise multiple comparisons (not assuming equal variances) was used to test differences among cytotypes. SPSS for Windows vs. 12 was used for all calculations.

RESULTS

Spatial distribution and frequencies of cytotypes

The geographical distribution of cytotypes largely confirms a spatial separation of the two most frequent ploidy levels, the diploids and tetraploids (Fig. 1). The distribution of diploids is confined to the south-western borders of the Alps outside the range of the last previous glacial maximum. Tetraploids are distributed all over the Alps in the previously glaciated area, including the population near the eastern margin of the former ice-shield. Also the geographically isolated populations at Mt Cusna and Corsica are tetraploid. A more diverse zone occurs at the overlapping distribution area of diploids and tetraploids in the south-western Alps, where diploids, triploids, tetraploids, pentaploids and hexaploids are sympatric, and co-occur sometimes in the same population. Notably, triploids and hexaploids also do occur occasionally outside the hybrid zone, in Gran Paradiso (no. 33) and in the Bernina massif (no. 47). Only diploid and tetraploid cytotypes are predominant within their respective populations (100% frequencies; Fig. 1). Triploid, pentaploid and hexaploid cytotypes are less stable, because they occur either only in mixed populations, or as low-frequency cytotypes within predominantly diploid or tetraploid populations. Within mixed populations, frequencies of triploids are highest among all the other rare cytotypes.

Mode of reproduction

FCSS revealed an unexpected high diversity of reproductive pathways (Table 3). FCSS confirms that diploids are sexual with a 2x embryo and a 3x endosperm. There was only one seed sample from a triploid mother plant (Allos, no. 23), which appeared to be apomictic and pseudogamous. Tetraploid mother plants showed a broader range of variation in seed formation, indicating that the species has not a fixed apomixis. The majority of tetraploid plants, are apomeiotic and pseudogamous with the use of one or two pollen nuclei for fertilization of the endosperm (4x embryos and 10x to 12x endosperm); two cases of autonomous endosperm formation (8x) are also indicated. Three individuals from the Col della Perla population (no. 12) had a fully sexual reproduction with 4x embryos and 6x endosperms. One tetraploid plant from Grand St Bernard (no. 35) had seeds with a hexaploid embryo, and a highly ploid, 20x endosperm. Here fertilization of an unreduced 4x egg cell by 2x pollen, and an endopolyploid endosperm is likely. Surprisingly, several samples showed triploid embryos as in Fig. 3. As they came from tetraploid parents, the embryo sac must have been formed via an unbalanced meiosis to produce triploid egg cells and polar nuclei. The egg cell developed parthenogenetically, while the endosperm was probably fertilized by one or two diploid sperm nuclei. Only this way can the ratio of 3x:8x or 3x:

10x be explained. These seeds occur in a couple of populations with predominantly 4x adult plants outside the putative hybrid zone. Several tetraploid plants formed 3x or 4x embryos, but no endosperm peak was detected. Here either abortion or a rapid consumption of the endosperm by the growing embryo could have happened, as it is known from Asteraceae (Krahulcova and Suda, 2006). Pentaploid individuals from the Col della Perla population (no. 11) seem to form reduced embryo sacs and fertilized egg cells. No seeds were available for analysis from hexaploid plants.

Fertility of cytotypes

The diploid cytotype has the highest fertility, with 30.8–97.1% of achenes within collective fruits well developed, while all the polyploids ranged from 0 to 60.9% (Table 4 and Fig. 4). The tetraploid cytotype falls with its maximum within the range of the diploids, but the interquartile range remains below the minimum of the diploids (Fig. 4). However, several samples (15 individuals) had no well-developed achenes at all, including the population from Mt Cusna. This explains that also the mean values of tetraploids remain low. ANOVA revealed a significant difference between percentages of well-developed achenes among the cytotypes (5x and 6x pooled; d.f. among groups = 3; d.f. within groups = 125; $F = 68.055$; $P < 0.001$). The pairwise multiple comparisons using Tanhame's *post hoc* tests revealed highly significant differences between the diploids and all polyploid cytotypes ($P < 0.001$ for all pairs). Between the 4x and the pooled 5x–6x cytotype, the difference is significant ($P = 0.017$), but there is neither a significant difference between 4x and 3x nor between 3x and 5x–6x cytotypes ($P > 0.5$).

DISCUSSION

Stability of tetraploid cytotypes enhances geographical parthenogenesis

The present data confirm that the diploid sexual populations are confined to a relic area in the south-western Alps outside the range of the ice cover of the last glacial maximum. On the whole distribution range in the previously glaciated area, tetraploids predominate; they are stable in the sense that only occasionally do other cytotypes occur within the tetraploid populations. Since tetraploids are confirmed to be predominantly apomictic or at least facultatively apomictic, these results confirm geographical parthenogenesis in this species.

Within the tetraploid populations, only a few rare hexaploid and triploid individuals occur far away from the putative hybrid zone where they are common. It seems that new cytotypes arise occasionally from partial apomixis (i.e. by uncoupling of apomeiosis and parthenogenesis), but they do have some difficulty in becoming established. They probably encounter few mating partners of the same ploidy level, resulting in meiotic disturbances and a lower fitness in the offspring. These factors may block the establishment of a critical number of individuals for survival of the new cytotype, as Levin (1975, 2002) described under the term 'minority cytotype disadvantage'. The apomictic tetraploid cytotype, in contrast,

TABLE 3. Summary of seed flow cytometric data with the observed ploidy levels in embryo and endosperm, and the inferred mode of reproduction

Population no.	Population name	Individual no.	Embryo	Second peak/G ₂ phase	Endosperm	Mode of reproduction			
						Origin of embryo sac	Egg cell	Endosperm development*	Mother plant [†]
8	Passo del Duca 9534	04	3x		8x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	NA
		08	3x		8x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	NA
11	Col della Perla I 9596	13	5x	10x	8x	Meiotic	Fertilized	Sexual	5x
12	Col della Perla II 9597	07	3x	6x	8x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x
		01	4x		6x	Meiotic	Fertilized	Sexual	4x
		03	4x	8x	6x	Meiotic	Fertilized	Sexual	4x
		03 (new)	4x	5x	6x	Meiotic	Fertilized	Sexual	4x
		08	4x		10x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	5x
		15	4x		10x	Apomeiotic	Parthenogenetic	Pseudogamous I	4x
		26	4x		8x	Apomeiotic	Parthenogenetic	Autonomous	4x
13	Notre Dame de la Fenestre	05	3x		10x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x
		02	4x		12x	Apomeiotic	Parthenogenetic	Pseudogamous II	4x
15	Colle della Lombarde I 9601	01	2x		3x	Meiotic	Fertilized	Sexual	2x
21	Cayolle II	09-1	4x		12x	Apomeiotic	Parthenogenetic	Pseudogamous II	4x
		09-2	4x		10x	Apomeiotic	Parthenogenetic	Pseudogamous I	4x
		15	4x		8x	Apomeiotic	Parthenogenetic	Autonomous	4x
23	Allos II	09	3x		8x	Apomeiotic	Parthenogenetic	Pseudogamous I	3x
24	Vars	06	3x		NA	Disturbed meiotic	Parthenogenetic	Aborted	4x
		15	3x		NA	Disturbed meiotic	Parthenogenetic	Aborted	4x
		16	3x		NA	Disturbed meiotic	Parthenogenetic	Aborted	4x
		05-1	2x		3x	Meiotic	Fertilized	Sexual	2x
29	Vercors II	05-2	2x		3x	Meiotic	Fertilized	Sexual	2x
		06	2x		3x	Meiotic	Fertilized	Sexual	2x
		07	2x		3x	Meiotic	Fertilized	Sexual	2x
		19	2x		3x	Meiotic	Fertilized	Sexual	2x
		23	2x		3x	Meiotic	Fertilized	Sexual	2x
		05	2x		3x	Meiotic	Fertilized	Sexual	2x
31	Mt Cenis	06	4x		NA	Apomeiotic	Parthenogenetic	Aborted	4x
32	Col d'Iseran	01	4x		NA	Apomeiotic	Parthenogenetic	Aborted	4x
		02	4x		NA	Apomeiotic	Parthenogenetic	Aborted	4x
		11	3x	5x	8x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x
33	Grand St Bernard	14	3x		NA	Disturbed meiotic	Parthenogenetic	Aborted	4x
		15	3x		14x	Disturbed meiotic	Parthenogenetic	Pseudogamous II	4x
		11 + 12	4x	6x	12x	Apomeiotic	Parthenogenetic	Pseudogamous II and sexual	4x
		12	4x		10x	Apomeiotic	Parthenogenetic	Pseudogamous I	4x
		13 (new)	4x		10x	Apomeiotic	Parthenogenetic	Pseudogamous I	4x
		21	4x		10x	Apomeiotic	Parthenogenetic	Pseudogamous I	4x
		21	4x		10x	Apomeiotic	Parthenogenetic	Pseudogamous I	4x
		26	4x		NA	Apomeiotic	Parthenogenetic	Aborted	4x
37	Ovronnaz	13	6x		20x	Apomeiotic	Fertilized	Pseudogamous II	4x
		19	4x		12x	Apomeiotic	Parthenogenetic	Pseudogamous II	4x
40	Lötschental	26	3x		8x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x
41	Simplon Pass	21	3x		10x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x
		25	3x	6x	9x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x

Continued

TABLE 3. *Continued*

Population no.	Population name	Individual no.	Embryo	Second peak/G ₂ phase	Endosperm	Mode of reproduction			Mother plant [†]
						Origin of embryo sac	Egg cell	Endosperm development*	
43	Lukmanier Pass	08	4x		12x	Apomeiotic	Parthenogenetic	Pseudogamous II	4x
		18	4x		NA	Apomeiotic	Parthenogenetic	Aborted	4x
		23	4x		12x	Apomeiotic	Parthenogenetic	Pseudogamous II	4x
		12	3x		8x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x
44	Rheinwald	26	3x		12x	Disturbed meiotic	Parthenogenetic	Pseudogamous II	4x
		03	3x		10x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x
47	Bernina Pass	01	3x		10x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x
		22	3x		8x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x
57	Mt Großglockner	02	3x		9x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x
		Herbar	3x		10x	Disturbed meiotic	Parthenogenetic	Pseudogamous II	4x
58	Mt Sadnig	Herbar	3x		8x	Disturbed meiotic	Parthenogenetic	Pseudogamous I	4x
		01	4x		10x	Apomeiotic	Parthenogenetic	Pseudogamous I	4x
59	Turracherhöhe	02	4x		10x	Apomeiotic	Parthenogenetic	Pseudogamous I	4x
		03	4x		10x	Apomeiotic	Parthenogenetic	Pseudogamous I	4x
		03	4x		10x	Apomeiotic	Parthenogenetic	Pseudogamous I	4x

NA, Not applicable.

* Pseudogamous I and II refer to a fertilization by one or two pollen nuclei, respectively (see also Hörandl *et al.*, 2008).

[†] Data are taken from leaf measurements.

TABLE 4. *Descriptive statistics of the percentages of well-developed achenes per collective fruit*

Cytotypes	<i>N</i>	Mean	Minimum	Maximum	s.d.
2x	21	67.79	30.77	97.14	17.96
3x	9	11.73	1.25	33.33	9.96
4x	93	17.96	0.00	60.87	15.51
5x	5	7.83	0.00	15.56	6.24
6x	1	3.17	3.17	3.17	

N = no. of collective fruits analysed.

avoids negative effects of meiotic disturbances and does not need a mating partner. Therefore, apomicts do not suffer from being a minority in the population (Hörandl, 2006), and can readily establish even aneuploid populations, as shown, for example, by Mraz *et al.* (2008) in *Pilosella*. For endosperm fertilization, apomictic *R. kuepferi* can use either self-pollen (Huber, 1988; E. Hörandl, unpubl. res.) or pollen of another cytotype in the population. The great variation in ploidy levels observed in the endosperm of well-developed seed suggests that seed formation in *R. kuepferi* is not highly sensitive against endosperm imbalance, similar to some pseudogamous Rosaceae (Talent and Dickinson, 2007). Therefore, pollination by another cytotype would not confer a minority cytotype disadvantage for endosperm formation in this species. Autonomous endosperm formation, as also observed occasionally in *R. kuepferi*, is completely pollen-independent and may enhance uniparental reproduction. Nevertheless,




FIG. 4. Boxplots of the variation of percentages of well-developed achenes per collective fruit for each cytotype. The box shows the 25th and 75th percentile range and the median value. *N* = number of collective fruits.

strong endosperm imbalance could potentially contribute to the high amount of seed abortion in polyploid cytotypes.

A different pathway may lead to the formation of pentaploids in the contact zone: one seed sample from a 5x mother plant indicates a fully sexual pathway, probably via a meiotic reduction of the embryo sac (3x) and fertilization by diploid pollen to form a 5x embryo and a 8x endosperm (Table 3). The origin of these pentaploids from other

cytotypes, however, may also involve apomeiotic and parthenogenetic pathways.

Outside of the contact zone, no diploids occur which would explain the formation of triploids. But, triploid embryos seem to arise quite often in the seed of tetraploid populations (Table 3 and Fig. 3), but almost never appear in the leaf material of adult plants. The reduction in ploidy level in the embryo can only be explained by a meiotically reduced embryo sac. It is supposed that meiosis was disturbed in the mother plant, resulting in megaspores with unbalanced chromosome numbers. Molecular markers (microsatellites, AFLPs) do not indicate the contribution of another parental species to the genome of the tetraploid cytotype (A.-C. Cosendai and E. Hörandl, unpubl. res.). Autopolyploid origin and multivalent formation during meiosis may cause such unbalanced chromosome numbers. The ratio obtained in the endosperm seems to confirm this. Effectively, with ratio $3x$ (embryo) : $8x$; $9x$; $10x$ (endosperm), the endosperm nuclei would be $6x$, plus $2x$, $3x$ and $4x$ derived from the pollen. Since the $3x$ embryo must have developed parthenogenetically, again either one or both pollen nuclei ($2x + 2x$) could have been used for endosperm fertilization. In some of the samples with $3x$ embryos (populations 12 and 41), an additional, but smaller, $6x$ peak was observed, which represents the G_2 peak of the growing embryo (e.g. Fig. 3). Moreover, disturbances in microsporogenesis and formation of unbalanced pollen (Izmailow, 1965, 1973, 1976; Jankun, 1965) can contribute to variation in the endosperm ploidy levels. Most importantly, these tetraploid plants obviously show only a partial apomixis by keeping a disturbed meiosis, but developing the egg cell parthenogenetically. In the next generations, further reduction of ploidy levels would finally lead to haploid offspring, in which recessive deleterious mutations would be fully expressed. These processes may strongly reduce fitness of such lineages with only a partial apomixis (e.g. Van Dijk and Vijverberg 2005). The present data show only a single triploid adult plant in the whole area (Fig. 1 and Table 2), but 17 triploid embryos in the seeds from tetraploid adults. These plants come from 11 populations outside the area of diploid influence, suggesting a rather frequent phenomenon (Table 3). However, triploids that have been formed via this partial apomixis probably have difficulty in becoming established. In contrast, the fully apomictic pathway with a combination of apomeiosis and parthenogenesis maintains the tetraploid level and is obviously more successful in the establishment of this cytotype, as seen in the frequency of adult plants. In two of the populations analysed (Grand St Bernard and Simplon Pass), both $3x$ and $4x$ embryos occur in the seeds of tetraploids, but both populations had 100% adult tetraploid plants. In populations with both pathways, selection obviously favours individuals expressing full apomixis, which stabilizes the ploidy level, over parthenogenesis alone which forms new cytotypes.

Hexaploids can be formed in a tetraploid population from a cross between an unreduced embryo and reduced pollen. In this case, the endosperm would be $10x$ ($8x + 2x$). Only one seed sample with a $6x$ embryo and an $20x$ endosperm were found. Here endopolyploidy could explain the double DNA content of the expected $10x$. Endopolyploidy in the endosperm is known, for example, from *Zea mays* and other species (Kowles *et al.*, 1988; Barow and Meister, 2003; Barow,

2006). Such hexaploid B_{III} hybrids may also give rise to triploids via meiosis and parthenogenetic development of the egg cell, but are obviously not stable.

These aspects would also explain why coexistence of cytotypes is rare in *R. kuepferi*, in contrast to what Kao (2007) showed for apomictic *Arnica cordifolia*. In *Arnica*, apomictic reproduction is predominant and probably stable enough to keep cytotypes reproductively isolated. Additionally, differences in phenology maintain a stable coexistence of cytotypes. Stable triploidy is also widespread in dandelions (*Taraxacum officinale* group; Van Dijk, 2003). In Asteraceae, apomixis may actually arise at the triploid level because of their mode of embryo sac development and endosperm formation (Talent, 2009). In *Ranunculus kuepferi*, the present results rather suggest that apomixis originated in tetraploid cytotypes, because of the existence of rare sexual tetraploids and a frequent uncoupling of apomeiosis and parthenogenesis on this ploidy level.

The variety of pathways indicates again the broad flexibility of modes of reproduction within this species. It appears to be able to express all possible solutions of embryo formation between fully sexual to complete apomixis. The coupling of apomeiosis and parthenogenetic development of the egg cell seems not yet fully established. Furthermore, *R. kuepferi* can undergo all kinds of endosperm development from the pseudogamous to the autonomous type. These phenomena suggest a very young, postglacial or even recurrent evolutionary origin of apomixis. Further support for this hypothesis comes from linear correlations of genome size to ploidy levels (Fig. 2), which infers that the frequently observed genome downsizing in polyploids has not yet occurred (see Leitch and Bennett, 2004).

Stability of the diploid sexual populations

Diploids maintain themselves via sexual reproduction and high seed set, but they can accept pollen from tetraploids in the sympatric zone to form triploid or pentaploid offspring. In the hybrid zone, triploids and pentaploids may be recurrently formed but still they are quite rare and do not form populations on their own. The relatively frequent triploids can arise from different pathways: a crossing between a haploid meiotic egg cell (x) and tetraploid meiotic pollen ($2x$) gives a triploid embryo, while the endosperm ($2x$) plus one pollen nucleus ($2x$) would be tetraploid; this $3x : 4x$ ratio was not actually observed in the wild seeds, but could account in for triploids being in the majority in the present data. The tetraploid pollen donor could be either sexual or apomictic. Alternatively, within a diploid population, occasionally unreduced pollen ($2x$) can be formed, which, when it fertilizes a haploid egg cell, results in the same embryo : endosperm ratio as above. The reciprocal cross would retrieve a $3x : 5x$ ratio. These processes could be a spontaneous step towards autopolyploidy via the triploid bridge (Ramsey and Schemske, 1998) and would not involve a cross between cytotypes. The present limited sampling of achenes in diploid populations may account for the lack of evidence on these pathways.

The hypothesis that apomixis can be transferred to sexual populations via the pollen of apomicts would suggest a

replacement of sexual by apomictic cytotypes (Mogie, 1992; Mogie *et al.*, 2007). There is little evidence for this pathway in the present dataset, partly because of low seed set in triploids. A triploid plant from the Allos population (no. 23), that shows fully apomictic reproduction (Table 3), may have indeed originated from an apomictic pollen donor. In the seed material, the number of triploid embryos in the hybrid zone is very low (three), but these embryos have originated from a tetraploid mother plant. Therefore, introgression of apomixis into diploid sexuals cannot be proven directly with the present seed dataset. Leaf material, however, shows that almost all triploid adult plants come from the hybrid zone, suggesting that crossings between diploid and tetraploid cytotypes probably do occur recurrently. This does not necessarily indicate an introgression process, because the tetraploid parents in this area could also be fully sexual, as shown in three individuals from the Col della Perla population (no. 12 in Table 3). If triploids in the hybrid zones could be sexual or partly sexual, then disturbances of meiosis, reduced fitness and minority cytotype disadvantages may strongly limit the establishment of triploid lineages. Triploid apomicts, in contrast, should be able to establish purely triploid populations, but such populations have not been observed. It is concluded that triploids originated mainly from sexual events rather than from triploid apomictic parents.

The present results indicate that introgression of apomixis into sexual populations in the $2x-4x$ hybrid zone may occur, but frequencies need to be studied further. It remains questionable whether this process plays a major role for the observed pattern of geographical parthenogenesis. Moreover, the diploid sexual individuals of *R. kuepferi* do have a significantly higher fertility than all the polyploid cytotypes (Fig. 4). In such cases, it is unlikely that apomixis can replace sexuality in mixed populations (see Mogie, 1992; Hörandl and Tensch, 2009). This is in accordance with results from *R. auricomus*, where findings of low crossability between different cytotypes, and a significantly higher fertility of sexuals rejected the introgression hypothesis (Hörandl and Tensch, 2009). Low frequencies of introgression were also found in population studies on *Taraxacum* (Brock, 2004).

Colonizing abilities of tetraploids

The easternmost population at Turracherhöhe and the outposts at Corsica and Mt Cusna are all tetraploid. For the population at Turracherhöhe, the FCSS data (Table 3) revealed fully apomictic reproduction. The samples from Mt Cusna had completely aborted achenes, as is typical for apomicts. There were no achenes in the populations in Corsica, but Huber (1989) also reported seed and pollen abortion. Apomictic reproduction in both the Corsica and Mt Cusna populations is further indicated by a clonal population genetic structure (Cosendai, 2005; unpubl. res.). These outposts have been most probably founded by efficiently colonizing tetraploid apomicts, probably via long-distance dispersal. Corsica has had no land bridge to the European continent since the Miocene (Loÿe-Pilot *et al.*, 2004). The location of Mt Cusna in the northern Apennines may lead to a hypothesis that the species could have migrated from the Alps towards the Apennines, but there is no high mountain in Liguria to support this idea.

As the diploid populations occupy a somewhat central position between Corsica and the major tetraploid area, it is likely that tetraploids originated in this area, but expanded their range rapidly via a centrifugal dispersal. Such distribution patterns of central diploid sexuals surrounded by expanding polyploid cytotype apomicts are common in other cases of geographical parthenogenesis [*Antennaria* (Bayer, 1990); *Stevia* (Soejima *et al.*, 2001); *Paspalum* (Urbani, 2002); *Taraxacum* and *Chondrilla* (Van Dijk, 2003); *Ranunculus auricomus* (Hörandl, 2009)]. Long-distance dispersal, even between continents, is a frequent phenomenon in the genus *Ranunculus* (K. Emadzade and E. Hörandl, unpubl. res.). The achenes of *R. kuepferi* are small and have a cavity between the seed and the pericarp, aiding wind dispersal (Müller-Schneider, 1986). Otherwise, birds or mammals may have been other important dispersal vectors. The ability of apomicts to found populations with a single diaspore may have enhanced the success of rare dispersal events (Hörandl *et al.*, 2008). Since the apomicts do not have a fitness advantage with respect to female fecundity, it is likely that superior colonizing abilities, as postulated by Baker's Law (Baker, 1965, 1967), are a main factor for the observed centrifugal distribution. For the diploid sexual cytotype, self-sterility (Huber, 1988) and pollinator-dependence may strongly limit range expansions.

Conclusions

The results on *R. kuepferi* overall confirm a high stability of the diploid and the tetraploid cytotype in their respective areas. Aneuploid cytotypes are probably recurrently formed, but fail to establish as long as remnants of the sexual pathway are maintained. The stability of the apomictic tetraploid cytotypes may consequently be important for establishing a large distribution area. In contrast, the significantly higher fertility of the diploids may stabilize the diploid cytotype in its relic area. Moreover, the present data tend to prove the difficulty for a new cytotype to establish, as long as partial sexuality is maintained; minority cytotype effects *sensu* Levin (1975), Husband (2000) and Husband *et al.* (2008) seem to play an important role in this pattern for reducing the presence of different cytotypes in the population, although the seeds are constantly newly produced.

ACKNOWLEDGEMENTS

Financial support from the Austrian Research Foundation (FWF, project P19006-B03) and the Austrian Academy of Sciences, Commission for Interdisciplinary Ecological studies (KIÖS, project P2009-03), both granted to E.H., is gratefully acknowledged. We thank Eva Maria Tensch and Johann Greilhuber (both at Vienna) for introducing us to flow cytometry and for helpful discussions, our colleagues for collection of materials, and two anonymous referees for valuable comments on the manuscript. Funding to pay the Open Access publication charges for this article was provided by the Austrian Research Foundation (FWF).

LITERATURE CITED

Asker SE, Jerling L. 1992. *Apomixis in plants*. Boca Raton, FL: CRC Press.

- Baker HG. 1965.** Characteristics and modes of origin of weeds. In: Baker HG, Stebbins GL. eds. *The genetics of colonizing species*. New York, NY: Academic Press.
- Baker HG. 1967.** Support for Baker's law – as a rule. *Evolution* **21**: 853–856.
- Baranyi M, Greilhuber J. 1996.** Flow cytometric and feulgen densitometric analysis of genome size variation in *Pisum*. *Theoretical and Applied Genetics* **92**: 297–307.
- Barow M. 2006.** Endopolyploidy in seed plants. *Bioessays* **28**: 271–281.
- Barow M, Meister A. 2003.** Endopolyploidy in seed plants is differently correlated to systematics, organ, life strategy and genome size. *Plant, Cell & Environment* **26**: 571–584.
- Bayer RJ. 1990.** Investigations into the evolutionary history of the *Antennaria rosea* (Asteraceae: Inuleae) polyploid complex. *Plant Systematics and Evolution* **169**: 97–110.
- Bell G. 1982.** *The masterpiece of nature: the evolution and genetics of sexuality*. Berkeley, CA: California Press.
- Bierzchudek P. 1985.** Patterns in plant parthenogenesis. *Experientia* **41**: 1255–1264.
- Brock MT. 2004.** The potential for genetic assimilation of a native dandelion species, *Taraxacum ceratophorum* (Asteraceae), by the exotic congener *T. officinale*. *American Journal of Botany* **91**: 656–663.
- Burnier J, Burki S, Arrigo N, Küpfer P, Alvarez N. 2009.** Genetic structure and evolution of alpine polyploid complexes: *Ranunculus kuepferi* (Ranunculaceae) as a case study. *Molecular Ecology* **18**: 3720–3744.
- Comai L. 2005.** The advantages and disadvantages of being polyploid. *Nature Reviews Genetics* **6**: 836–846.
- Cosendai A-C. 2005.** *Liens de parenté et origine de Ranunculus kuepferi en Corse et dans le sud de la France*. MSc Thesis, University of Neuchâtel, Neuchâtel.
- Dickinson TA, Lo E, Talent N. 2007.** Polyploidy, reproductive biology, and Rosaceae: understanding evolution and making classifications. *Plant Systematics and Evolution* **266**: 59–78.
- Doležel J, Greilhuber J, Lucretti S, et al. 1998.** Plant genome size estimation by flow cytometry: inter-laboratory comparison. *Annals of Botany* **82** (Suppl. A): 17–26.
- Doležel J, Greilhuber J, Suda J. 2007a.** Estimation of nuclear DNA content in plants using flow cytometry. *Nature Protocols* **2**: 2233–2244.
- Doležel J, Greilhuber J, Suda J. 2007b.** *Flow cytometry with plant cells: analysis of genes, chromosomes and genomes*. Weinheim: Wiley-VCH.
- Galbraith DW, Harkins KR, Maddox JM, Ayres NM, Sharma DP, Firoozabady E. 1983.** Rapid flow cytometric analysis of the cell cycle in intact plant tissues. *Science* **220**: 1049–1051.
- Greilhuber J. 2008.** Cytochemistry and C-values: the less-well-known world of nuclear DNA amounts. *Annals of Botany* **101**: 791–804.
- Greilhuber J, Lysák MA, Doležel J, Bennett MD. 2005.** The origin, evolution and proposed stabilisation of the terms 'genome size' and 'C-value' to describe nuclear DNA contents. *Annals of Botany* **94**: 255–260.
- Hörandl E. 2006.** The complex causality of geographical parthenogenesis. *New Phytologist* **171**: 525–538.
- Hörandl E. 2008.** Evolutionary implications of self-compatibility and reproductive fitness in the apomictic *Ranunculus auricomus* polyploid complex (Ranunculaceae). *International Journal of Plant Sciences* **169**: 1219–1228.
- Hörandl E. 2009a.** Geographical parthenogenesis: opportunities for asexuality. In: Schoen I, Martens K, Van Dijk P. eds. *Lost sex*. Heidelberg: Springer, 161–186.
- Hörandl E. 2009b.** The evolution of self-fertility in apomictic plants. *Sexual Plant Reproduction* DOI 10.1007/s00497-009-0122-3.
- Hörandl E, Tensch EM. 2009.** Introgression of apomixis into sexual species is inhibited by mentor effects and ploidy barriers in the *Ranunculus auricomus* complex. *Annals of Botany* **104**: 81–89.
- Hörandl E, Cosendai AC, Tensch EM. 2008.** Understanding the geographic distributions of apomictic plants: a case for a pluralistic approach. *Plant Ecology and Diversity* **1**: 309–320.
- Huber W. 1988.** Natürliche Bastardierungen zwischen weißblühenden *Ranunculus*-Arten in den Alpen [Natural hybridizations between white-flowered species of *Ranunculus* in the Alps] [in German with English abstract]. *Veröffentlichungen des Geobotanischen Institutes der ETH Zürich* **100**: 1–160.
- Huber W. 1989.** *Ranunculus kuepferi* Greuter & Burdet in Korsica (Gruppe *R. pyrenaicus* L.). *Candollea* **44**: 630–637.
- Husband BC. 2000.** Constraints on polyploid evolution: a test of the minority cytotype exclusion principle. *Proceedings of the Royal Society of London Series B – Biological Sciences* **267**: 217–223.
- Husband BC, Ozimec B, Martin SL, Pollock L. 2008.** Mating consequences of polyploid evolution in flowering plants: current trends and insights from synthetic polyploids. *International Journal of Plant Sciences* **169**: 195–206.
- Izmailov R. 1965.** Macrosporogenesis in apomictic species *Ranunculus casubicus*. *Acta Biologica Cracoviensia Series Botanica* **8**: 183–195.
- Izmailov R. 1973.** Cyto-embryological studies in experimental hybrids of apomictic species *Ranunculus-cassubicus* L. *Acta Biologica Cracoviensia Series Botanica* **16**: 99–120.
- Izmailov R. 1976.** Problem of apomixis in *Ranunculus-auricomus* group. *Acta Biologica Cracoviensia Series Botanica* **19**: 15–28.
- Jankun A. 1965.** Studies of meiosis in various chromosomal types of *Ranunculus casubicus* L. *Acta Biologica Cracoviensia Series Botanica* **8**: 171–180.
- Kao RH. 2007.** Asexuality and the coexistence of cytotypes. *New Phytologist* **175**: 764–772.
- Kearney M. 2005.** Hybridization, glaciation and geographical parthenogenesis. *Trends in Ecology & Evolution* **20**: 495–502.
- Kearney M. 2006.** Response to Lundmark: polyploidization, hybridization and geographical parthenogenesis. *Trends in Ecology & Evolution* **21**: 10.
- Kowles RV, Phillips RL. 1988.** Endosperm development in maize. *International Review of Cytology* **112**: 97–136.
- Krahulcova A, Suda J. 2006.** A modified method of flow cytometric seed screen simplifies the quantification of progeny classes with different ploidy levels. *Biologia Plantarum* **50**: 457–460.
- Küpfer P. 1974.** The affinities between orophyte flora of the Alps and that of the Pyrenees. *Boissiera* **23**: 1–322.
- Law JH, Crespi BJ. 2002.** The evolution of geographic parthenogenesis in timema walking-sticks. *Molecular Ecology* **11**: 1471–1489.
- Leitch IJ, Bennett MD. 2004.** Genome downsizing in polyploid plants. *Biological Journal of the Linnean Society* **82**: 651–663.
- Levin DA. 1975.** Somatic cell hybridization application in plant systematics. *Taxon* **24**: 261–270.
- Levin DA. 2002.** *The role of chromosomal change in plant evolution*. Oxford: Oxford University Press.
- Loye-Pilot MD, Durand-Delga M, Feinberg H, Gourinard Y, Magne J. 2004.** The Burdigalian of eastern Corsica within its geodynamic setting. *Comptes Rendus Geoscience* **336**: 919–930.
- Lundmark M. 2006.** Polyploidization, hybridization and geographical parthenogenesis. *Trends in Ecology & Evolution* **21**: 9.
- Lynch M. 1984.** Destabilizing hybridization, general-purpose genotypes and geographic parthenogenesis. *Quarterly Review of Biology* **59**: 257–290.
- Matzk F, Meister A, Schubert I. 2000.** An efficient screen for reproductive pathways using mature seeds of monocots and dicots. *The Plant Journal* **21**: 97–108.
- Merxmüller H. 1952.** Untersuchungen zur Sipplgliederung und Arealbildung in den Alpen. Teil 1. *Jahrbuch des Vereins zum Schutze der Alpenpflanzen und -tiere* **17**: 96–133.
- Merxmüller H. 1953.** Untersuchungen zur Sipplgliederung und Arealbildung in den Alpen. Teil 2. *Jahrbuch des Vereins zum Schutze der Alpenpflanzen und -tiere* **18**: 135–158.
- Merxmüller H. 1954.** Untersuchungen zur Sipplgliederung und Arealbildung in den Alpen. Teil 3. *Jahrbuch des Vereins zum Schutze der Alpenpflanzen und -tiere* **19**: 97–139.
- Mogie M. 1992.** *The evolution of asexual reproduction in plants*. London: Chapman and Hall.
- Mogie M, Britton NF, Stewart-Cox JA. 2007.** Asexuality, polyploidy and the male function. In: Hörandl E, Grossniklaus U, Van Dijk PJ, Sharbel T. eds. *Apomixis: evolution, mechanisms and perspectives*. Ruggell, Liechtenstein: ARG-Gantner, 195–214.
- Mraz P, Singliarova B, Urfus T, Krahulec F. 2008.** Cytoecography of *Pilosella officinarum* (Compositae): altitudinal and longitudinal differences in ploidy level distribution in the Czech republic and Slovakia and the general pattern in Europe. *Annals of Botany* **101**: 59–71.
- Müller-Schneider P. 1986.** Verbreitungsbiologie der Blütenpflanzen Graubündens [Diasporology of the spermatophytes of the Grisons (Switzerland)]. *Berichte des Geobotanischen Institutes ETH, Zürich* **85**: 1–263.
- Nogler GA. 1984a.** Gametophytic apomixis. Johri BM. ed. *Embryology of angiosperms*. Berlin: Springer-Verlag, 475–566.

- Nogler GA. 1984b.** Genetics of apospory in apomictic *Ranunculus auricomus*. 5. Conclusion. *Botanica Helvetica* **94**: 411–422.
- Nogler GA. 1995.** Genetics of apomixis in *Ranunculus auricomus*. 6. Epilogue. *Botanica Helvetica* **105**: 111–115.
- Parisod C, Besnard G. 2007.** Glacial *in situ* survival in the western Alps and polytopic autopolyploidy in *Biscutella laevigata* L. (Brassicaceae). *Molecular Ecology* **16**: 2755–2767.
- Paun O, Stuessy TF, Hörandl E. 2006.** The role of hybridization, polyploidization and glaciation in the origin and evolution of the apomictic *Ranunculus cassubicus* complex. *New Phytologist* **171**: 223–236.
- Ramsey J, Schemske DW. 1998.** Pathways, mechanisms, and rates of polyploid formation in flowering plants. *Annual Review of Ecology and Systematics* **29**: 467–501.
- Schönswetter P, Stehlik I, Holderegger R, Tribsch A. 2005.** Molecular evidence for glacial refugia of mountain plants in the European Alps. *Molecular Ecology* **14**: 3547–3555.
- Soejima A, Yahara T, Watanabe K. 2001.** Distribution and variation of sexual and agamospermous populations of *Stevia* (Asteraceae: Eupatorieae) in the lower latitudes, Mexico. *Plant Species Biology* **16**: 91–105.
- Spielman M, Vinkenoog R, Scott RJ. 2003.** Genetic mechanisms of apomixis. *Philosophical Transactions of the Royal Society of London Series B – Biological Sciences* **358**: 1095–1103.
- Suda J, Trávníček P. 2006.** Estimation of relative nuclear DNA content in dehydrated plant tissues by flow cytometry. *Current Protocols in Cytometry* Chapter 7: Unit 7.30.
- Talent N. 2009.** Evolution of gametophytic apomixis in flowering plants: an alternative model from maloid Rosaceae. *Theory in Biosciences* **128**: 121–138.
- Talent N, Dickinson TA. 2007.** Endosperm formation in aposporous *Crataegus* (Rosaceae, Spiraeoideae, tribe Pyreae): parallels to Ranunculaceae and Poaceae. *New Phytologist* **173**: 231–249.
- Temsch EM, Greilhuber J, Hammett KRW, Murray BG. 2008.** Genome size in *Dahlia* cav. (Asteraceae-Coreopsidae). *Plant Systematics and Evolution* **276**: 157–166.
- Temsch EM, Greilhuber J, Krisai R. 2009.** Genome size in liverworts. *Preslia*, in press.
- Thompson SL, Whitton J. 2006.** Patterns of recurrent evolution and geographic parthenogenesis within apomictic polyploid Easter daises (*Townsendia hookeri*). *Molecular Ecology* **15**: 3389–3400.
- Tribsch A. 2004.** Areas of endemism of vascular plants in the eastern Alps in relation to Pleistocene glaciation. *Journal of Biogeography* **31**: 747–760.
- Urbani MH, Quarin CL, Espinoza F, Penteadó MIO, Rodrigues IF. 2002.** Cyto geography and reproduction of the *Paspalum simplex* polyploid complex. *Plant Systematics and Evolution* **236**: 99–105.
- Van Dijk PJ. 2003.** Ecological and evolutionary opportunities of apomixis: insights from *Taraxacum* and *Chondrilla*. *Philosophical Transactions of the Royal Society of London Series B – Biological Sciences* **358**: 1113–1120.
- Van Dijk PJ. 2007.** Potential and realized costs of sex in dandelions, *Taraxacum officinale* s.l. In: Hörandl E, Grossniklaus U, Van Dijk PJ, Sharbel T. eds. *Apomixis: evolution, mechanisms and perspectives*. Ruggell, Liechtenstein: ARG-Gantner, 215–233.
- Van Dijk PJ, Vijverberg K. 2005.** The significance of apomixis in the evolution of the angiosperms: a reappraisal. In: Bakker F, Chatrou L, Gravendeel B, Pelsers PB. eds. *Plant species-level systematics: new perspectives on pattern and process*. Ruggell, Liechtenstein: Gantner, 101–116.
- Vandel A. 1928.** La parthénogenèse géographique. Contribution à l'étude biologique et cytologique de la parthénogenèse naturelle. *Bulletin Biologique de la France et de la Belgique* **62**: 164–182.
- Van Valen L. 1973.** A new evolutionary law. *Evolutionary Theory* **1**: 1–30.
- Vinkenoog R, Bushell C, Spielmann M, Adams S, Dickinson HG, Scott RJ. 2003.** Genomic imprinting and endosperm development in flowering plants. *Molecular Biotechnology* **25**: 149–184.
- Vorburger C. 2006.** Geographic parthenogenesis: recurrent patterns down under. *Current Biology* **16**: R641–R643.
- Vrijenhoek RC. 1984.** Ecological differentiation among clones: the frozen niche variation model. In: Woermann K, Loeschcke V. eds. *Population biology and evolution*. Berlin: Springer, 217–231.
- Vrijenhoek RC. 1994.** Unisexual fish: Model systems for studying ecology and evolution. *Annual Review of Ecology and Systematics* **25**: 71–96.
- Vuille C, Küpfer P. 1985.** Aposporie chez le *Ranunculus parnassifolius*. L. I. Etude cytoembryologique. *Bulletin de la Société Neuchâteloise des Sciences Naturelles* **108**: 123–134.