

## Population dynamics of epiphytic orchids in a metapopulation context

Manuela Winkler<sup>1,\*</sup>, Karl Hülber<sup>2</sup> and Peter Hietz<sup>1</sup>

<sup>1</sup>Institute of Botany, Department of Integrative Biology, University of Natural Resources and Applied Life Sciences (Boku), Gregor-Mendel-Str. 33, 1180 Vienna, Austria and <sup>2</sup>VINCA – Vienna Institute for Nature Conservation & Analyses, Giessergasse 6/7, 1090 Vienna, Austria

Received: 7 April 2009 Returned for revision: 2 June 2009 Accepted: 29 June 2009 Published electronically: 11 August 2009

- **Background and Aims** Populations of many epiphytes show a patchy distribution where clusters of plants growing on individual trees are spatially separated and may thus function as metapopulations. Seed dispersal is necessary to (re)colonize unoccupied habitats, and to transfer seeds from high- to low-competition patches. Increasing dispersal distances, however, reduces local fecundity and the probability that seeds will find a safe site outside the original patch. Thus, there is a conflict between seed survival and colonization.
- **Methods** Populations of three epiphytic orchids were monitored over three years in a Mexican humid montane forest and analysed with spatially averaged and with spatially explicit matrix metapopulation models. In the latter, population dynamics at the scale of the subpopulations (epiphytes on individual host trees) are based on detailed stage-structured observations of transition probabilities and trees are connected by a dispersal function.
- **Key Results** Population growth rates differed among trees and years. While ignoring these differences, and averaging the population matrices over trees, yields negative population growth, metapopulation models predict stable or growing populations because the trees that support growing subpopulations determine the growth of the metapopulation. Stochastic models which account for the differences among years differed only marginally from deterministic models. Population growth rates were significantly lower, and extinctions of local patches more frequent in models where higher dispersal results in reduced local fecundity compared with hypothetical models where this is not the case. The difference between the two models increased with increasing mean dispersal distance. Though recolonization events increased with dispersal distance, this could not compensate the losses due to reduced local fecundity.
- **Conclusions** For epiphytes, metapopulation models are useful to capture processes beyond the level of the single host tree, but local processes are equally important to understand epiphyte population dynamics.

**Key words:** Dispersal, epiphyte, fecundity, *Jacquiella leucomelana*, *Jacquiella teretifolia*, *Lycaste aromatica*, matrix population model, metapopulation model, population dynamics, conflict between seed survival and colonization.

### INTRODUCTION

The spatial distribution of individuals in a plant population is mainly determined by seed dispersal patterns and subsequent establishment success. Most wind-dispersed seeds land near their source and very few travel over large distances (Bullock and Clarke, 2000; Nathan and Muller-Landau, 2000; Tackenberg, 2003). Seedling establishment in orchids can be facilitated by conspecific plants because the chance of forming associations with beneficial mycorrhizal fungi is higher close to established plants (Diez, 2007). However, with increasing plant density, fecundity and survival in a population decrease as a result of increasing competition (e.g. Watkinson and Harper, 1978; Mithen *et al.*, 1984), or as a result of increased attraction of herbivores and seed predators (Feeny, 1976). Dispersal, in turn, is not only influenced by species-specific seed characteristics, but also by the distribution of adult plants and local demographic processes, primarily fecundity (Clark and Ji, 1995; Levin *et al.*, 2003).

In a metapopulation, seed dispersal is necessary to colonize unoccupied habitats, to re-colonize extinct patches, and to transfer seeds from high to low competition patches (Husband and Barrett, 1996). However, high dispersal distances reduce local fecundity and the probability that seeds will reach a suitable habitat. Thus, there is a conflict between seed survival and colonization (Levin *et al.*, 2003), the existence of which was demonstrated for a hypothetical metapopulation (Johst *et al.*, 2002) and for *Pinus halepensis* (Bohrer *et al.*, 2005). Whether long-distance dispersal is beneficial depends on local population dynamics. For instance, long-range dispersal has a positive effect on metapopulation persistence unless the number of potential dispersers is low due to small population growth rates (Johst *et al.*, 2002). The interactions between fecundity and mean dispersal distance, and the consequences for population maintenance, are little studied.

The effect of spatial distribution in populations is now an essential component of metapopulation models, and has become firmly established in population biology (Hanski and Simberloff, 1997). Classical metapopulation theory focuses

\* For correspondence. E-mail manuela.winkler@boku.ac.at

on the extinction and re-colonization of local patches regarding neither size and spatial arrangement of the patches nor local population dynamics (Levins, 1969). More recently, metapopulation theory has been considerably extended to include spatially explicit and realistic approaches including declining non-equilibrium, habitat-tracking, and mainland-island metapopulations (Thomas, 1994; Hanski, 1997; Harrison and Taylor, 1997). Most metapopulation models, however, concentrate on regional dynamics and ignore details on the scale of local populations (but see Bohrer *et al.*, 2005; Volis *et al.*, 2005; Mildén *et al.*, 2006).

Previous studies on the population dynamics of vascular epiphytes found survival to be the most important parameter determining population growth rates (Hernández-Apolinar, 1992; Zotz *et al.*, 2005; Zotz and Schmidt, 2006; Winkler *et al.*, 2007), but none of these studies accounted for the spatial structure of epiphyte populations. In the present study, it is demonstrated how the incorporation of spatial structure alters model predictions of population size and the importance of demographic processes for population growth rates of three epiphytic orchids. Spatially realistic matrix metapopulation models are used, where population dynamics at the scale of the local population (individuals growing on one host tree) are based on detailed stage-structured observations of transition probabilities and epiphyte populations on different trees are connected by a dispersal function. The question is asked whether demographic processes differ significantly between trees, and if so, what the consequences for metapopulation structure and persistence are. Furthermore, metapopulations where increased dispersal results in reduced local fecundity are compared with hypothetical models where local fecundity is not related to dispersal.

## METHODS AND RESULTS

### Study area and species

This study was conducted in the forest reserve ‘Santuario Bosque de Niebla’ adjacent to the Instituto de Ecología, 2.5 km south of Xalapa, in central Veracruz, Mexico (19°31'N, 96°57'W; 1350 m a.s.l.). Average temperature is 19 °C, and annual precipitation is approx. 1500 mm, most of which falls in the rainy season (June to October). The forest is commonly classified as ‘bosque mesófilo de montaña’ (mesophilous montane forest) following Rzedowski (1986). A description of the epiphyte community is given by Hietz and Hietz-Seifert (1995).

The epiphytic orchids *Jacquinella leucomelana* (Reichenbach f.) Schlechter, *J. teretifolia* (Sw.) Britton & P. Wilson, and *Lycaste aromatica* (Graham ex Hook.) Lindley were censused. Reproductive plants of *J. leucomelana* and *J. teretifolia* were found on trees with a diameter at breast height (dbh) >14 cm, and *L. aromatica* only on trees >50 cm dbh (Hietz *et al.*, 2006). None of the species is restricted to a single host tree species although there are preferences for *Quercus* spp., which are dominating the forest (Hietz *et al.*, 2006).

### Population census

In an area of approx. 3 ha within the Santuario Bosque de Niebla all accessible individuals on ten trees were tagged in August 2001. Individuals on a tree were treated as a subpopulation, resulting in five subpopulations of *J. leucomelana*, and six of *J. teretifolia* and *L. aromatica* (Fig. 1). According to epiphyte density estimates in this part of the forest, approx. 25 % of *J. leucomelana*, 60 % of *J. teretifolia*, and 75 % of *L. aromatica* individuals were sampled (Hietz *et al.*, 2006; M. Winkler, unpubl. res.). In February of the years 2002–2005, survival, vegetative growth and reproduction of the tagged individuals, and new seedlings were recorded. In the *Jacquinella* species, the ramets, open flowers, dehiscent flowers and fruits were counted, and the length of the longest ramet was measured to the nearest 0.5 cm. In *L. aromatica*, the number of peduncles and fruits was recorded. The height, breadth and width of each living pseudobulb of *L. aromatica* were measured in the first census, and only height of the new pseudobulb in the subsequent censuses because width and length are highly correlated with height (Winkler, 2001). Pseudobulb volume was calculated as an ellipsoid

$$V_p = (2\pi/3)HBW \quad (1)$$

where *H*, *B* and *W* are pseudobulb height, breadth and width, respectively. The sum of the volume of all living pseudobulbs was the size of an individual.

### Spatially averaged matrix population models

To characterize population dynamics, the stage-classified transition matrices of Lefkovich (1965) were used. Four stage classes were distinguished with a combination of reproductive and size criteria (Table 1). The size threshold of seedlings (*s*) was defined as the 95th percentile size of 1-year-old or younger seedlings. Non-reproductive adults (*a*) were considered to be at least the size of the 5th percentile of reproductive plants, i.e. plants that had reached the size of reproducing individuals (*f*), but had not yet produced flowers or currently had none. Other individuals were grouped as juveniles (*j*).

Transition matrices (*A*), where the individuals of all host trees were pooled, were built for each of the three projection intervals (February 2002 to February 2003, 2003–2004, and 2004–2005), and a mean matrix was calculated using the function ‘mean.list’ of the R-library popbio (R 2.7.0; R Development Core Team, 2008; Stubben and Milligan, 2007). These matrices contain the probabilities of transition from stage *x* at the beginning of a projection interval (a year, time *t* – 1) to stage *y* at the end of the interval (time *t*). Fecundity (*F*, i.e. the ‘transition’ from *f*-stages to the *s*-stage) is the number of newborn seedlings (*n<sub>s</sub>*) at time *t* divided by the number of reproductive individuals (*n<sub>f</sub>*) at time *t* – 1. Deterministic population growth rates  $\lambda_{det}$  (the dominant eigenvalue of the transition matrix) were calculated following Caswell (2001) for single-year matrices (see Table S1 in Supplementary Data, available online) and for mean matrices (Table 2).




FIG. 1. Characteristics of epiphyte subpopulations on ten trees: position of occupied trees, seed flow as estimated with a dispersal model for orchid seeds (Murren and Ellison, 1998), and number of reproductive individuals. Arrows indicate the direction and magnitude of seed flow between trees with numbers indicating the mean probability (%) over three years (February in years 2002–2005) that a seedling on a target tree originates from a given source tree. The difference between the indicated probability and 100 % equates to seeds originating from the same tree (population). Trees not connected by arrows contribute <0.1 % of seeds. The size of the tree symbol is proportional to the tree diameter.

TABLE 1. Definition of stage classes according to the presence of flowers or fruits (F), length of the longest ramet (L), number of pseudobulbs ( $N_p$ ) and cumulative pseudobulb volume ( $V_p$ )

Species	Criterion	Stage			
		Seedlings (s)	Juveniles (j)	Non-reproductive adults (a)	Reproductive adults (f)
<i>Jacquinella leucomelana</i>	F	No	No	No	Yes
	L (cm)	≤1.5	>1.5–3	>3	Any
<i>J. teretifolia</i>	F	No	No	No	Yes
	L (cm)	≤1.5	>1.5–10.5	>10.5	Any
<i>Lycaste aromatica</i>	F	No	No	No	Yes
	$N_p$	1	Any	Any	Any
	$V_p$ (cm <sup>3</sup> )	≤0.4	>0.4–31.5	>31.5	Any

For each mean matrix, an elasticity matrix was constructed: between trees (Caswell, 2001):

$$e_{ij} = \left( \frac{a_{ij}}{\lambda} \frac{\delta \lambda}{\delta a_{ij}} \right) \quad (2)$$

$$\mathbf{A}_m = \begin{pmatrix} \mathbf{A}_{\text{tree 1}} & \mathbf{M}_{\text{tree 1} \rightarrow 2} & \dots & \mathbf{M}_{\text{tree 1} \rightarrow n} \\ \mathbf{M}_{\text{tree 2} \rightarrow 1} & \mathbf{A}_{\text{tree 2}} & \dots & \mathbf{M}_{\text{tree 2} \rightarrow n} \\ \dots & \dots & \dots & \dots \\ \mathbf{M}_{\text{tree } n \rightarrow 1} & \mathbf{M}_{\text{tree } n \rightarrow 2} & \dots & \mathbf{A}_{\text{tree } n} \end{pmatrix} \quad (3)$$

where  $a_{ij}$  is an entry in the transition matrix. The elasticity value for a particular matrix element ( $e_{ij}$ ) measures the effect of changes in  $a_{ij}$  on  $\lambda$  (Caswell, 2001). Elasticity values were summed up across demographic processes (survival, growth, retrogression, fecundity and dispersal) to represent their proportional impact on population growth rates (Silvertown *et al.*, 1993; Fig. 3A).

Unless stated otherwise, all analyses were computed using S-Plus 2000 (MathSoft Inc., 1988–1999).

#### Metapopulation matrix models

To account for the spatial structure of the populations, metapopulation matrices ( $\mathbf{A}_m$ ) were constructed (in the following ‘empirical metapopulation models’). These matrices consist of tree-specific transition submatrices ( $\mathbf{A}_{\text{tree } 1 \dots n}$ ) on the main diagonal and migration submatrices ( $\mathbf{M}$ ) describing dispersal

M-submatrices contain only zeros except for fecundity. In metapopulation models the fecundity ( $F_m$ ) value in each sub-matrix represents the number of newborn seedlings on a target tree ( $\text{tree}_{\text{TA}}$ ) divided by the number of reproductive plants of a source tree ( $\text{tree}_{\text{SO}}$ ):

$$F_m = \frac{n_{s\text{-SO} \rightarrow \text{TA}}}{n_{f\text{-SO}}} \quad (4)$$

The seedlings on each  $\text{tree}_{\text{TA}}$  were assigned to a source tree proportional to the number of seeds that tree contributed to the seed rain reaching a target tree assuming equal germination and survival of seeds from all source trees.

$$n_{s\text{-SO} \rightarrow \text{TA}} = \frac{n_{\text{seeds-SO} \rightarrow \text{TA}}}{n_{\text{seeds-TOTAL}}} \quad (5)$$

TABLE 2. Comparison of deterministic ( $\lambda_{\text{det}}$ ) and stochastic ( $\lambda_s$ ) growth rates of populations of epiphytic orchids

	SUM	MM <sub>emp</sub>	MM <sub>adj</sub>
<i>Jacquinella leucomelana</i>			
$\lambda_{\text{det}}$	0.895	1.025	0.986
$\lambda_s$ (95 % CI)	0.893 (0.892–0.894)	1.025 (1.024–1.025)	0.963 (0.960–0.965)
<i>J. teretifolia</i>			
$\lambda_{\text{det}}$	0.980	1.127	1.106
$\lambda_s$ (95 % CI)	0.960 (0.959–0.962)	1.110 (1.109–1.111)	1.089 (1.086–1.091)
<i>Lycaste aromatica</i>			
$\lambda_{\text{det}}$	0.949	1.007	1.007
$\lambda_s$ (95 % CI)	0.949 (0.948–0.949)	1.007 (1.006–1.007)	1.007 (1.006–1.007)

Given are  $\lambda_{\text{det}}$  of mean matrices over three years and  $\lambda_s$  (with 95 % confidence interval from simulations over 50 000 time steps) for spatially averaged matrix models (SUM) and metapopulation models [MM<sub>emp</sub> = metapopulation models where the transitions within each host tree and dispersal between host trees are calculated separately (empirical metapopulation model); MM<sub>adj</sub> = metapopulation models based on MM<sub>emp</sub> where transition probabilities for stages with an initial frequency of less than five individuals were replaced by probabilities of the corresponding spatially averaged matrix model (adjusted metapopulation model)].




FIG. 2. Calculation of the probability  $p_{\text{SO} \rightarrow \text{TA}}$  that a seed produced by epiphytic orchids on a source tree ( $\text{tree}_{\text{SO}}$ ) was dispersed to a target tree ( $\text{tree}_{\text{TA}}$ ). This probability corresponds to the area under the dispersal kernel  $k$  covered by the target tree (hatched area) in relation to the total area covered by the seed shadow of the source tree (area enclosed by the outermost ring). The shaded area is the area at the distance  $d_{\text{SO} \rightarrow \text{TA}} \pm c_{\text{TA}}/2$  (crown diameter of  $\text{tree}_{\text{TA}}$ ) between  $\text{tree}_{\text{SO}}$  and  $\text{tree}_{\text{TA}}$  (compare eqns 8 and 9 in the text).

As seed rain is difficult to measure in the field, the number of reproductive plants on  $\text{tree}_{\text{SO}}$  ( $n_{\text{f-SO}}$ ) was used, and the total number of reproductive plants ( $n_{\text{f-TOTAL}}$ ) as a proxy for their

seed yield assuming equal seed output per reproductive plant.

$$n_{\text{seeds-TOTAL}} \approx n_{\text{f-TOTAL}} \quad (6)$$

$$n_{\text{seeds-SO} \rightarrow \text{TA}} \approx n_{\text{f-SO}} \times p_{\text{SO} \rightarrow \text{TA}} \quad (7)$$

where  $p_{\text{SO} \rightarrow \text{TA}}$  is the probability that a seed produced on  $\text{tree}_{\text{SO}}$  was dispersed to  $\text{tree}_{\text{TA}}$ , assuming equal dispersal in all directions. This probability is reflected by the area under the dispersal kernel  $k$  at the distance  $d_{\text{SO} \rightarrow \text{TA}} \pm c_{\text{TA}}/2$  (crown diameter of  $\text{tree}_{\text{TA}}$ ) between  $\text{tree}_{\text{SO}}$  and  $\text{tree}_{\text{TA}}$  in relation to total curve area (Fig. 2):

$$p_{\text{SO} \rightarrow \text{TA}} = \frac{c_{\text{TA}}}{2d_{\text{SO} \rightarrow \text{TA}} \times \pi} \times \int_{d_{\text{SO} \rightarrow \text{TA}} - \frac{c_{\text{TA}}}{2}}^{d_{\text{SO} \rightarrow \text{TA}} + \frac{c_{\text{TA}}}{2}} \int_0^\infty k \quad (8)$$

$$\text{if } \text{tree}_{\text{SO}} \neq \text{tree}_{\text{TA}} \quad p_{\text{SO} \rightarrow \text{TA}} = \frac{c_{\text{TA}}}{2} \int_0^\infty \int_0^\infty k \quad \text{if } \text{tree}_{\text{SO}} = \text{tree}_{\text{TA}} \quad (9)$$

where  $k$  is the distance-probability function describing the dispersal kernel (following Clark *et al.*, 1998), a function of the distance between trees ( $d_{\text{SO} \rightarrow \text{TA}}$ ) and the mean dispersal distance of seeds ( $X$ ):

$$k = e^{-\frac{d_{\text{SO} \rightarrow \text{TA}}}{X}} \quad (10)$$

Mean dispersal distance of seeds depends on seed and canopy characteristics, and on release height, and was calculated following Murren and Ellison (1998):

$$X = \frac{U_f e^{\frac{\alpha h - z}{\lambda}}}{\lambda U_t} \quad (11)$$

where  $U_f$  (wind velocity above canopy,  $4.9 \text{ m s}^{-1}$ ),  $\alpha$  (degree of canopy openness, 2–14),  $\lambda$  (factor describing the spread of settling particles around their estimated mean location, 1.53), and  $U_t$  (terminal seed velocity,  $0.157 \text{ m s}^{-1}$ ) were adopted from Murren and Ellison's model for the epiphytic orchid *Brassavola nodosa* (Murren and Ellison, 1998). The mean canopy height in the forest ( $z$ ) is 24 m (Williams-Linera and Tolome, 1996), and  $h$ , the seed release height, was set to the






FIG. 3. Elasticity of the (meta)population growth rate to demographic processes (survival, growth, retrogression, fecundity, dispersal) of epiphytic orchids. The elasticity values from mean matrices are given over three years (February in 2002–2005): a spatially averaged matrix model (SUM); a metapopulation model where the transitions within each host tree and dispersal between host trees are calculated separately (empirical metapopulation model,  $MM_{emp}$ ); and metapopulation models where fecundity ( $F$ ) and mean dispersal distance ( $X$ ) were each multiplied by 0.5, 1 and 2 to create nine scenarios each for models including a conflict between seed survival and colonization (open symbols), and for models lacking this conflict (i.e. within-tree fecundity constant regardless of dispersal; closed symbols).

mean height of reproductive plants in a given source tree at the time of a given census. Empirical metapopulation models for each of the three years are shown in Table S2 in Supplementary Data, available online.

Because transition probabilities become inaccurate with low population sizes, an ‘adjusted metapopulation model’ was also calculated where transition probabilities for stages with an initial frequency of less than five individuals were replaced by probabilities of the spatially averaged matrix model, i.e. the mean of the entire population. Deterministic growth rates of mean empirical and adjusted metapopulation matrices are given in Table 2 and elasticity values in Fig. 3A.

#### Spatially averaged matrix model vs. metapopulation model

Deterministic population growth rates of mean spatially averaged matrices were below one, and of empirical metapopulations above one in all species. Population growth rates of adjusted metapopulation models, where transition probabilities of a small subpopulation are replaced by the population mean, were also consistently higher than those of spatially averaged matrices (Table 2).

In both spatially averaged and metapopulation models, survival was the most important demographic process as indicated by elasticity analysis (Fig. 3). In the *Jacquiniella* species, the elasticity of population growth rate to survival increased when incorporating spatial structure, whereas the elasticity to growth, retrogression and fecundity decreased. In *Lycaste aromatica*, this pattern was reversed. Elasticity analysis of the empirical metapopulation model showed that dispersal had no effect on metapopulation growth rate in any of the species studied.

#### Effect of temporal and spatial variation

To evaluate the effect of temporal (observation years) and spatial (host trees representing subpopulations) variation in transition probabilities on the fate of a plant, loglinear analyses were carried out following Horvitz and Schemske (1995) and Caswell (2001). The dependence of individual fate ( $F$ ) on the explanatory variables state ( $S$ ), year ( $Y$ ) and tree ( $T$ ) based on four-way contingency tables derived from transition probabilities of the empirical metapopulation model was examined. It was assumed that  $F$  is affected by  $S$ . Thus, all models included the  $SF$  interaction and the three-way interaction of explanatory variables, entailing the null model  $STY,SF$ . Additional models included the interactions of  $F$  with  $Y$ ,  $T$  or both, leading to the models  $STY,SFY$ ,  $STY,SFT$  and  $STY,SFY,SFT$ . These models were compared to the saturated model ( $SFYT$ ), which considered all possible interactions. The model with the lowest Akaike information criterion (AIC) was regarded as the best model. The AIC-value of the best model was subtracted from that of the other models; rescaled values are given as  $\Delta AIC$  (Burnham and Anderson 1998; Caswell, 2001).  $STY,SFY,SFT$  (i.e. interactions of  $F$  with both  $Y$  and  $T$ ) was the best model for both *Jacquiniella* species. *Jacquiniella leucomelana* was more variable among years (i.e. compared with  $STY,SF$  the model  $STY,SFY$  improved the model fit more than  $STY,SFT$ ), while *J. teretifolia* showed stronger variability across host trees. For *L. aromatica* neither temporal nor spatial variation was found (Table 3).

TABLE 3. Log-linear analyses of the effect of the host tree (T) and year (Y) on fates (F) of epiphytic orchids conditional on their initial state (S) based on four-way contingency tables (T,Y,S,F) following Caswell (2001)

Model	Log-likelihood ratio	d.f.	P	AIC	ΔAIC
<i>Jacquinella leucomelana</i>					
STY,SF	581.1	176	<0.001	229.1	276.5
STY,SFY	251.8	144	<0.001	-36.2	11.2
STY,SFT	473.4	128	<0.001	217.4	264.8
<b>STY,SFY,SFT</b>	144.6	96	0.001	-47.4	0
SFYT	0	0		0	47.4
<i>Jacquinella teretifolia</i>					
STY,SF	682.9	272	<0.001	138.9	397.9
STY,SFY	441.4	240	<0.001	-38.6	220.4
STY,SFT	245.7	192	0.005	-138.3	120.8
<b>STY,SFY,SFT</b>	60.9	160	1.000	-259.1	0
SFYT	0	0		0	259.1
<i>Lycaste aromatica</i>					
<b>STY,SF</b>	336.1	236	<0.001	-135.9	0
STY,SFY	281.0	204	<0.001	-127.1	8.8
STY,SFT	189.5	156	0.035	-122.5	13.4
STY,SFY,SFT	138.8	124	0.172	-109.2	26.7
SFYT	0	0		72*	207.9

P-values indicate the significance of the deviation of a model from the saturated model SFYT. The model with the lowest Akaike information criterion (AIC) was regarded as the best model (given in bold). Log-likelihood ratio statistic was used to calculate AIC.

\* AIC of the saturated model deviates from zero, because one degree of freedom will be deducted for each structural zero by function loglm (Venables and Ripley, 2002).

To account for environmental stochasticity, the environment was modelled using independent and identically distributed sequences (Caswell 2001). This sequence of independent and identically distributed random variables was used to select one of the three yearly matrices. Resulting stochastic population growth rates ( $\lambda_s$ , calculated using the function 'stoch.-growth.rate' of the R-library popbio with 50 000 time intervals) were similar to deterministic growth rates both for spatially averaged, and empirical and adjusted metapopulation models (Table 2). The empirical metapopulation model showed that two subpopulations of *J. teretifolia*, and one of *J. leucomelana* and *L. aromatica*, respectively, were growing or stable (data not shown). The number of extinctions (a tree was scored as 'occupied' if it harboured at least one orchid individual in > 50 % of 1000 repetitions) and subsequent recolonizations in the course of 50 years was recorded. The number of occupied trees decreased to two in *J. leucomelana*, and three in *J. teretifolia* and *L. aromatica* after 50 years. The empirical metapopulation model predicted no recolonization of extinct subpopulations in any of the species (data not shown).

#### Fecundity, dispersal distance and the conflict between seed survival and colonization

In an epiphyte metapopulation, a seed can either stay within its natal host tree or be dispersed beyond it (conservation-of-individuals law), in which case seed survival is less probable and local fecundity is reduced. In other words, a larger mean dispersal distance ( $X$ ) in eqn (11) results in a 'fatter' tail of the dispersal kernel, but simultaneously fewer seeds are left within the crown of the source tree. Thus local fecundity and population growth rate will decrease with increasing  $X$ . To demonstrate the effects of this conflict on metapopulation growth rates, and extinction/recolonization

dynamics, a series of metapopulation scenarios was constructed with varying fecundities ( $F$ ) and mean dispersal distances ( $X$ ) based on transition probabilities of the adjusted metapopulation models. Fecundity in the transition ( $A_{tree1 \dots n}$ ) and migration ( $M$ ) submatrices was calculated as:

$$F_m = Fmp_{SO \rightarrow TA} \quad (12)$$

where  $F$  is the fecundity of the corresponding spatially averaged matrix ( $A$ ), and  $m$  is a multiplication vector (containing 0.5, 1 and 2),  $p_{SO \rightarrow TA}$  was calculated applying eqns (8)–(11), where  $X$  was increased or decreased relative to the original value (i.e. multiplied with  $m$ ). A combination of the variations in  $X$  and  $F$  results in nine scenarios ( $F0.5-X0.5$ ,  $F0.5-X1$ ,  $F0.5-X2$ ,  $F1-X0.5$ ,  $F1-X1$ ,  $F1-X2$ ,  $F2-X0.5$ ,  $F2-X1$  and  $F2-X2$ ). These scenarios were compared with hypothetical metapopulation models where the conservation-of-individuals law was invalidated, i.e. assuming no reduction of local fecundity with increasing  $X$  by keeping the number of seeds dispersed within  $tree_{SO}$  constant and independent of  $X$  (i.e.  $p_{SO \rightarrow TA} = 1$  for  $A_{tree1 \dots n}$  submatrices in eqn 12).

The difference in metapopulation growth rates between models with and without the conflict between seed dispersal and colonization increased with both increasing  $F$  and  $X$  (Fig. 4). Deterministic and stochastic metapopulation growth rates (calculated as described above) hardly differed, thus only  $\lambda_s$  values are shown. Elasticity analysis of mean matrices (carried out as described above) of both *Jacquinella* species indicated that, in metapopulations, models including the conflict between seed survival and colonization, survival and retrogression were more important for metapopulation growth rates compared with the models with the conservation-of-individuals law invalidated, while growth and fecundity were less important (Fig. 3B). The differences increased with increasing  $X$  but were not significant (pairwise




FIG. 4. Conflict between seed survival and colonization of epiphytic orchids demonstrated as the difference in stochastic population growth rates ( $\lambda_s$ ) between metapopulation matrix models with or without this conflict, as indicated. Given are simulated  $\lambda_s$  over 50 000 time steps ( $\pm 95\%$  CI). Three annual metapopulation transition matrices (February in years 2002–2005) where the transitions within each host tree and dispersal between host trees are calculated separately. Within-tree fecundity values ( $F$ ) and mean dispersal distance ( $X$ ) were multiplied by 0.5, 1 and 2 to create nine scenarios ( $F0-5-X0-5$ ,  $F0-5-X1$ ,  $F0-5-X2$ ,  $F1-X0-5$ ,  $F1-X1$ ,  $F1-X2$ ,  $F2-X0-5$ ,  $F2-X1$  and  $F2-X2$ ).

$t$ -test, all  $P > 0.05$ ). In *L. aromatica*, there were no differences between the two models.

Stochastic models were calculated for each of the scenarios as described above. At each time step, one of the three yearly matrices was randomly drawn with equal probability and multiplied with the population vector of the previous year. This process was repeated 1000 times. For each tree, extinction (i.e. no individuals left on a tree) and recolonization events in the course of 50 years in percentage of possible extinctions (tree is occupied at time  $t - 1$ ) and recolonizations events in

percentage of possible extinctions (tree is empty at time  $t - 1$ ), respectively, were recorded. Metapopulation models including the conflict between seed survival and colonization predicted more extinctions (Fig. 5A–C) and fewer recolonizations (Fig. 5D) than models with no such conflict. The larger  $X$  was within a given  $F$ , the stronger was the conflict. The difference in extinctions of *L. aromatica* and of recolonizations of *J. teretifolia* also increased with increasing  $F$ . Both  $F$  and  $X$  had a significant effect in all species (significant  $F:X$  interaction in two-factor ANOVAs, all  $P < 0.001$ ).

## DISCUSSION

### Spatial arrangement and model predictions


The populations of most epiphytic orchids studied with spatially averaged matrix models seem to be declining (Calvo, 1993; Tremblay and Ackerman, 2001; Zotz and Schmidt, 2006). Also, repeated photographs of epiphytes on eight trees revealed a negative balance between birth and mortality for *Encyclia ochracea*, *Jacquiniella teretifolia* and *Lycaste aromatica*, with only the creeping orchid *Dichaea neglecta* increasing (Hietz, 1997). Ignoring the differences among host trees and averaging matrices over trees, population growth rates below one suggest that the populations of the three epiphytic orchids studied were also declining (Table 2).

In contrast, Tremblay *et al.* (2006) applied Levin's metapopulation model (Levin, 1969) to analyse population dynamics of an epiphytic/lithophytic orchid alongside a riverbed and found that the metapopulation of *Lepanthes rupestris* is stable. However, their study did not account for local population dynamics and assumed all subpopulations to be equidistant and of equal size. Metapopulation models which relax unrealistic assumptions (Lopez and Pfister, 2001; Higgins and Cain, 2002; Baguette and Schtickzelle, 2003) suggest that neither spatial heterogeneity nor local population dynamics should be disregarded. Therefore, our present spatially realistic simulation approach (Hanski, 1997) combines classical population models not accounting for spatial structure at all and metapopulation models that entirely disregard within-tree (i.e. local patch) dynamics, enabling a direct comparison between models including and those ignoring spatial structure. In contrast to spatially averaged matrix models, metapopulation models predicted stable or even growing populations of the three orchids; also when transition probabilities were adjusted to account for low individual numbers on individual trees (Table 2).

### The role of trees in metapopulation dynamics

In both spatially averaged and metapopulation model survival is by far the most important demographic process determining population growth, followed by growth (Fig. 3A). Dispersal had next to no influence in the metapopulation model, thus local processes dominate over regional ones (compare, for example, Volis *et al.*, 2005). Why then do the population growth rates of the models diverge?

The fate of individuals of both *Jacquiniella* species was significantly affected by the host tree to which they were attached (Table 3; compare Horvitz and Schemske, 1995; Zotz and



Schmidt, 2006). This may be related to the size or species of the host tree. For instance, trees with larger branches and a more corrugated bark or thick bryophyte mats may offer better conditions to epiphytes in terms of water supply (Hietz *et al.*, 2002). Some of the *Jacquiniella* host trees are deciduous and the increased light availability compared with evergreen trees might be beneficial.

In the metapopulation models, only some trees had an influence on population growth rates. Whether a subpopulation contributed to the growth of the metapopulation was determined by its growth rate (Table S2 in Supplementary Data, available online). A subpopulation with a growth rate  $>1$  is predicted to grow and to persist, regardless of its initial size. In contrast, in a spatially averaged matrix where individuals from growing and shrinking subpopulations are lumped together, large subpopulations have more weight because they harbour more individuals. If these large subpopulations happen to be in decline, so is the whole population. Thus, spatially averaged models of epiphytic orchids might underestimate population growth, which could explain the negative growth rates found in most models of epiphytes populations (Calvo, 1993; Tremblay and Ackerman, 2001; Zotz and Schmidt, 2006).

#### Dispersal, fecundity and the conflict between seed survival and colonization

Dispersal is a key process that shapes the spatial dynamics of plant populations and influences the probabilities of colonizing new sites as well as the extinction of existing populations (Nathan and Muller-Landau, 2000). According to the present elasticity analysis, dispersal had only a very small effect on population growth rates (Fig. 3B). Although there was some seedling ‘exchange’ between trees, dispersal beyond adjacent trees was a very rare event in our models (Fig. 1). The leptokurtic dispersal kernel for wind-dispersed seeds based on wind-channel experiments (Murren and Ellison, 1998) used in the present study could have underestimated the ‘fatness’ of the tail of the curve (Bullock and Clarke, 2000). However, independent population genetic studies of orchids (Trapnell and Hamrick, 2004; Jacquemyn *et al.*, 2007; but see Trapnell *et al.*, 2004) also show that most recruitments occur within a few metres from the mother plant. Seed numbers in orchids are very high (Arditti and Ghani, 2000), reflecting the small chance of finding a safe site including a fungal partner for germination. The number of seedlings

FIG. 5. Conflict between seed survival and colonization of epiphytic orchids demonstrated as the difference in extinction and recolonization events between metapopulation matrix models with or without this conflict. Differences (mean  $\pm$  s.e.) are shown for (A–C) extinction events in percentage of possible extinctions (tree is occupied at time  $t - 1$ ) and (D) recolonization events in percentage of possible recolonizations (tree is empty at time  $t - 1$ ). Models were run over 50 years with 1000 repetitions of permutations of three annual transition matrices (February in years 2002–2005) where the transitions within each host tree and dispersal between host trees are calculated separately. Within-tree fecundity ( $F$ ) and mean dispersal distance ( $X$ ) were multiplied by 0.5, 1 and 2 to create nine scenarios. There were no recolonizations in *Lycaste aromatica* and  $<10$  in *Jacquiniella leucomelana*. Numbers indicate extinction and recolonization events (%) in the models without the conflict. \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$  ( $t$ -tests between models including and lacking the conflict).



relative to seed production (number of seeds/capsule  $\times$  number of fruits; M. Winkler and P. Hietz, unpubl.res.) was 0.4‰ in *J. leucomelana*, 0.1‰ in *J. teretifolia*, and only 0.006‰ in *L. aromatica*. Seed dispersal to other trees was therefore orders of magnitudes higher than effective seedling exchange. One caveat of the present metapopulation model is that it ignores colonization of new trees (Snäll *et al.*, 2003). *Lepanthes rupestris* initially colonizes unoccupied trees with a mean probability of 1.1 % per year (Tremblay *et al.* 2006), but distances between trees are much smaller than in the forest studied here. Thus, colonization events are probably very rare in the present system, given the low seedling exchange rates among local populations (Fig. 1).

As the chance of finding a safe site beyond the patch of origin is low, increasing dispersal distance results in reduced seed survival at the (meta)population level (Levin *et al.*, 2003; Bohrer *et al.*, 2005; Cousens *et al.*, 2008). The more seeds are dispersed beyond their natal tree and the farther they travel on average, the more get lost in the unsuitable matrix between host trees (e.g. Bohrer *et al.*, 2005) and, as they become unavailable for local fecundity, they reduce local population growth. This conflict was evident in the populations of the epiphytic orchids studied where population growth rates (Fig. 4), as well as extinctions and recolonizations of local patches (Fig. 5) differed significantly between models where local fecundity was affected by dispersal distance and those where local fecundity was kept constant.

Bohrer *et al.* (2005) suggest that selection should favour long-distance dispersal under intermediate local extinction frequencies. Populations with low extinction probabilities do not profit from long-distance dispersal because patch turnover is slow and most available patches are eventually occupied. In populations with high local extinction pressure, a high proportion of seeds lost through long-distance dispersal, most of which will not found a new population, would increase the probability of local extinction. Therefore traits enhancing local survival, including lower dispersal distance, should be selected for (Bohrer *et al.*, 2005). Apart from extinctions as a consequence of demographic stochasticity (overall low population size in combination with spatial isolation), epiphyte populations are particularly prone to environmental stochasticity such as branchfall (Hietz, 1997; Zotz *et al.*, 2005; Zotz and Schmidt, 2006), suggesting rather high local extinction rates. Particularly in species growing in large clusters, such as the orchids studied here, the fall of individual branches may wipe out the entire population on a tree, but the effect of high rates of extinctions caused by the host tree's death (Overton, 1994; Snäll *et al.*, 2005; Tremblay *et al.*, 2006) in this context remains unknown. With increasing mean dispersal distance extinction probabilities increased (Fig. 5A–C) and population growth rates decreased (Fig. 4) in the epiphytic orchids because more seeds travelled beyond the original – suitable – patch and were mostly lost. Though increasing mean dispersal distance also increased the number of recolonization events (Fig. 5D), these could not compensate for the reduced local fecundity (Fig. 4).

The conflict between seed survival and colonization was more pronounced in the two *Jacquiiniella* species, which prefer thinner branches and where survival had less and fecundity had more influence on population growth rates

than in *Lycaste* (Figs 4 and 5). In the latter species, where population growth rates are almost independent of fecundity, how many seeds get lost in the unsuitable matrix between host trees has little effect as long as the trees do not die.

### Conclusions

When analysing the population dynamics of three epiphytic orchid species in a metapopulation context, both regional (between-tree) and local (within-tree) processes were important in the populations of *Jacquiiniella* spp., though the latter dominated. Since most trees in the forest appear to be suitable habitats for *Jacquiiniella* spp. (in the sense that at least some individuals can germinate and establish there) they may be characterized as 'spatially extended populations' (Freckleton and Watkinson, 2002). However, if a host tree effect (e.g. availability of mycorrhizal fungi) considerably shortens the number of suitable trees, the populations may be categorized as 'regional ensembles' where the amount of unsuitable habitat exceeds that of suitable sites (Freckleton and Watkinson, 2002). In contrast to *Jacquiiniella* spp., the persistence of the non-equilibrium metapopulation (*sensu* Harrison and Taylor, 1997) in *L. aromatica* depended on a single subpopulation with measurable exchange only between two neighbouring trees (Table S2 in Supplementary Data available online). We found the metapopulation concept useful to capture processes beyond the level of a single host tree. It may produce more realistic and accurate analyses of epiphyte populations and uncover potential bias in spatially averaged matrix models. However, the analysis of the conflict between seed survival and colonization showed that processes ensuring local survival should be selected for, whereas long-distance dispersal played a minor role in population maintenance. Thus, also in metapopulation models, local demographic processes are important and need to be included.

### SUPPLEMENTARY DATA

Supplementary data are available online at [www.aob.oxfordjournals.org](http://www.aob.oxfordjournals.org) and consist of the following files. Table S1: Spatially averaged transition matrices of populations of epiphytic orchids. Table S2: Metapopulation matrices (empirical metapopulation models) of populations of epiphytic orchids.

### ACKNOWLEDGEMENTS

We are very grateful for Hal Caswell's constructive criticism. Pieter Zuidema and two anonymous reviewers made useful suggestions on a previous draft, and Charles Nock kindly edited the manuscript. We are indebted to the Instituto de Ecología in Xalapa, to José García Franco for general support, and to Leticia Cruz Paredes for help in the field. This work was supported by the Austrian Science Fund (FWF; grant numbers P14775, P17875).

### LITERATURE CITED

- Arditti J, Ghani AK. 2000. Numerical and physical properties of orchid seeds and their biological implications. Tansley Review No 110. *New Phytologist* 145: 367–421.

- Baguette M, Schtickzelle N. 2003. Local population dynamics are important to the conservation of metapopulations in highly fragmented landscapes. *Journal of Applied Ecology* 40: 404–412.
- Bohrer G, Nathan R, Volis S. 2005. Effects of long-distance dispersal for metapopulation survival and genetic structure at ecological time and spatial scales. *Journal of Ecology* 93: 1029–1040.
- Bullock JM, Clarke RT. 2000. Long distance seed dispersal by wind: measuring the tail of the curve. *Oecologia* 124: 506–521.
- Burnham KP, Anderson DR. 1998. *Model selection and inference: a practical information-theoretic approach*. Heidelberg: Springer.
- Calvo RN. 1993. Evolutionary demography of orchids: intensity and frequency of pollination and the cost of fruiting. *Ecology* 74: 1033–1042.
- Caswell H. 2001. *Matrix population models: construction, analysis, and interpretation*, 2nd edn. Sunderland, MA: Sinauer.
- Clark JS, Ji Y. 1995. Fecundity and dispersal in plant populations: implications for structure and diversity. *The American Naturalist* 146: 72–111.
- Clark JS, Macklin E, Wood L. 1998. Stages and spatial scales of recruitment limitation in southern Appalachian forests. *Ecological Monographs* 68: 213–235.
- Cousens R, Dytham C, Law R. 2008. *Dispersal in plants: a population perspective*. Oxford: Oxford University Press.
- Diez JM. 2007. Hierarchical patterns of symbiotic orchid germination linked to adult proximity and environmental gradients. *Journal of Ecology* 95: 159–170.
- Feeny PP. 1976. Plant apparency and chemical defense. In: Wallace J, Mansell RL eds. *Biochemical interactions between plants and insects*. Recent Advances in Phytochemistry Vol. 10. New York, NY: Plenum Press, 1–40.
- Freckleton RP, Watkinson AR. 2002. Large-scale spatial dynamics of plants: metapopulations, regional ensembles and patchy populations. *Journal of Ecology* 90: 419–434.
- Hanski I. 1997. Metapopulation dynamics: from concepts and observations to predictive models. In: Hanski I, Gilpin ME. eds. *Metapopulation biology: ecology, genetics and evolution*. San Diego, CA: Academic Press, 69–91.
- Hanski I, Simberloff D. 1997. The metapopulation approach, its history, conceptual domain, and application to conservation. In: Hanski I, Gilpin ME. eds. *Metapopulation biology: ecology, genetics and evolution*. San Diego, CA: Academic Press, 5–26.
- Harrison S, Taylor AD. 1997. Empirical evidence for metapopulation dynamics. In: Hanski I, Gilpin ME. eds. *Metapopulation biology: ecology, genetics and evolution*. San Diego, CA: Academic Press, 27–42.
- Hernández-Apolinar M. 1992. *Dinámica poblacional de Laelia speciosa (H.B.K.) Schltr. (Orchidaceae)*. Tesis de Licenciatura, Universidad Autónoma de México, México.
- Hietz P. 1997. Population dynamics of epiphytes in a Mexican humid montane forest. *Journal of Ecology* 85: 767–775.
- Hietz P, Hietz-Seifert U. 1995. Intra- and interspecific relations within an epiphyte community in a Mexican humid montane forest. *Selbyana* 16: 135–140.
- Hietz P, Wanek W, Wania R, Nadkarni NM. 2002. Nitrogen-15 natural abundance in a montane cloud forest canopy as an indicator of nitrogen cycling and epiphyte nutrition. *Oecologia* 131: 350–355.
- Hietz P, Buchberger G, Winkler M. 2006. Effect of forest disturbance on abundance and distribution of epiphytic bromeliads and orchids. *Ecotropica* 12: 103–112.
- Higgins SI, Cain ML. 2002. Spatially realistic plant metapopulation models and the colonisation-competition trade-off. *Journal of Ecology* 90: 616–626.
- Horvitz CC, Schemske DW. 1995. Spatiotemporal variation in demographic transitions of a tropical understory herb: projection matrix analysis. *Ecological Monographs* 65: 155–192.
- Husband BC, Barrett SC. 1996. A metapopulation perspective in plant population biology. *Journal of Ecology* 84: 461–469.
- Jacquemyn H, Brys R, Vandepitte K, Honnay O, Roldán-Ruiz I, Wiegand T. 2007. A spatially explicit analysis of seedling recruitment in the terrestrial orchid *Orchis purpurea*. *New Phytologist* 176: 448–459.
- Johst K, Brandl R, Eber S. 2002. Metapopulation persistence in dynamic landscapes: the role of dispersal distance. *Oikos* 98: 263–270.
- Lefkovich LP. 1965. The study of population growth in organisms grouped by stages. *Biometrics* 21: 1–18.
- Levin SA, Muller-Landau HC, Nathan R, Chave J. 2003. The ecology and evolution of seed dispersal: a theoretical perspective. *Annual Review of Ecology, Evolution and Systematics* 34: 575–604.
- Levins R. 1969. Some demographic and genetic consequences of environmental heterogeneity for biological control. *Bulletin of the Entomological Society of America* 15: 237–240.
- Lopez JE, Pfister CA. 2001. Local population dynamics in metapopulation models: implications for conservation. *Conservation Biology* 15: 1700–1709.
- Mildén M, Münzbergová Z, Herbén T, Ehrlén J. 2006. Metapopulation dynamics of a perennial plant, *Succisa pratensis*, in an agricultural landscape. *Ecological Modelling* 199: 464–475.
- Mithen R, Harper JL, Weiner J. 1984. Growth and mortality of individual plants as a function of “available area”. *Oecologia* 62: 57–60.
- Murren CJ, Ellison AM. 1998. Effects of habitat, plant size, and floral display on male and female reproductive success of the Neotropical orchid *Brassavola nodosa*. *Biotropica* 28: 30–41.
- Nathan R, Muller-Landau HC. 2000. Spatial patterns of seed dispersal, their determinants and consequences for recruitment. *Trends in Ecology and Evolution* 15: 278–285.
- Overton JM. 1994. Dispersal and infection in mistletoe metapopulations. *Journal of Ecology* 82: 711–723.
- R Development Core Team. 2008. *R: a language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>.
- Rzedowski J. 1986. *Vegetación de México*, 3rd edn. Mexico: Editorial Limusa.
- Silvertown J, Franco M, Pisanty I, Mendoza A. 1993. Comparative plant demography: relative importance of life-cycle components to the finite rate of increase in woody and herbaceous perennials. *Journal of Ecology* 81: 465–467.
- Snäll T, Ribeiro PJ, Rydin H. 2003. Spatial occurrence and colonisations in patch-tracking metapopulations: local conditions versus dispersal. *Oikos* 103: 566–578.
- Snäll T, Ehrlén J, Rydin H. 2005. Colonisation-extinction dynamics of an epiphyte metapopulation in a dynamic landscape. *Ecology* 86: 106–115.
- Stubben CJ, Milligan BG. 2007. Estimating and analyzing demographic models using the popbio package in R. *Journal of Statistical Software* 22: 11.
- Tackenberg O. 2003. Modeling long-distance dispersal of plant diaspores by wind. *Ecological Monographs* 73: 173–189.
- Thomas CD. 1994. Extinction, colonisation, and metapopulations: environmental tracking by rare species. *Conservation Biology* 8: 373–378.
- Trapnell DW, Hamrick JL. 2004. Partitioning nuclear and chloroplast variation at multiple spatial scales in the neotropical epiphytic orchid, *Laelia rubescens*. *Molecular Ecology* 13: 2655–2666.
- Trapnell DW, Hamrick JL, Nason JD. 2004. Three-dimensional fine-scale genetic structure of the neotropical epiphytic orchid, *Laelia rubescens*. *Molecular Ecology* 13: 1111–1118.
- Tremblay RL, Ackerman JD. 2001. Gene flow and effective population size in *Lepanthes* (Orchidaceae): a case for genetic drift. *Biological Journal of the Linnean Society* 72: 47–62.
- Tremblay RL, Meléndez-Ackerman E, Kapan D. 2006. Do epiphytic orchids behave as metapopulations? Evidence from colonisation, extinction rates and asynchronous population dynamics. *Biological Conservation* 129: 70–81.
- Venables WN, Ripley BD. 2002. *Modern applied statistics with S*, 4th edn. New York, NY: Springer.
- Volis S, Bohrer G, Oostermeijer G, van Tienderen P. 2005. Regional consequences of local population demography and genetics in relation to habitat management in *Gentiana pneumonanthe*. *Conservation Biology* 19: 357–367.
- Watkinson AR, Harper JL. 1978. The demography of a sand dune annual: *Vulpia fasciculata*. I. The natural regulation of populations. *Journal of Ecology* 66: 15–33.
- Williams-Linera G, Tolome J. 1996. Litterfall, temperate and tropical dominant trees, and climate in a Mexican lower montane forest. *Biotropica* 28: 649–656.
- Winkler M. 2001. *Demographie der epiphytischen Orchideen* *Lycaste aromatica*, *Jacquinella leucomelana* und *J. teretifolia*. Diploma thesis, University of Vienna, Austria.
- Winkler M, Hülber K, Hietz P. 2007. Population dynamics of epiphytic bromeliads: life strategies and the role of host branches. *Basic and Applied Ecology* 8: 183–196.
- Zotz G, Schmidt G. 2006. Population decline in the epiphytic orchid *Aspasia principissa*. *Biological Conservation* 129: 82–90.
- Zotz G, Laube S, Schmidt G. 2005. Long-term population dynamics of the epiphytic bromeliad, *Werauhia sanguinolenta*. *Ecography* 28: 806–814.